arXiv:1509.03197v3 [math-ph] 18 Nov 2016

Superradiance initiated inside the ergoregion

Gregory Eskin,
Department of Mathematics, UCLA,
Los Angeles, CA 90095-1555, USA.
eskin@math.ucla.edu

November 21, 2016

Abstract

We consider the stationary metrics that have both the black hole and the ergoregion. The class of such metric contains, in particular, the Kerr metric. We study the Cauchy problem with highly oscillatory initial data supported in a neighborhood inside the ergoregion with some initial energy E_0. We prove that when the time variable x_0 increases this solution splits into two parts: one with the negative energy $-E_1$ ending at the event horizon in a finite time, and the second part, with the energy $E_2 = E_0 + E_1 > E_0$, escaping, under some conditions, to the infinity when $x_0 \rightarrow +\infty$. Thus we get the superradiance phenomenon. In the case of the Kerr metric the superradiance phenomenon is "short-lived", since both the solutions with positive and negative energies cross the outer event horizon in a finite time (modulo $O(\frac{1}{k})$) where k is a large parameter. We show that these solutions end on the singularity ring in a finite time. We study also the case of naked singularity.

Keywords: suprradiance, black holes, geometric optics
Mathematics Subject Classification 2010: 83C57, 83C75, 76J20

1 Introduction.

Consider the Lorentzian metric in $\mathbb{R}^3 \times \mathbb{R}$ of the form

$$dx_0^2 - \sum_{k=1}^{3} dx_k^2 - K(x)\Big(- dx_0 + \sum_{k=1}^{3} b_k(x) dx_k \Big)^2, \tag{1.1}$$

where $K(x) > 0,\ \sum_j^3 b_j^2(x) = 1$.

The ergoregion for this metric is the region where $K > 1$.

Let (ρ, φ, z) be the cylindrical coordinates in $\mathbb{R}^3$, i.e. $x_1 = \rho \cos \varphi,\ x_2 = \rho \sin \varphi,\ x_3 = z$, let x_0 be the time variable.

Write the inverse to the metric tensor in the cylindrical coordinates. Then the Hamiltonian $H(\rho, \varphi, z, \xi_0, \xi_\rho, \rho_\varphi, \xi_z)$ has the form

$$H = \xi_0^2 - \xi_\rho^2 - \frac{1}{\rho^2}\xi_\varphi^2 - \xi_z^2 + K(-\xi_0 + \hat{b} \cdot \hat{\xi})^2, \tag{1.2}$$

where $\hat{\xi} = (\xi_\rho, \frac{\xi_\varphi}{\rho}, \xi_z), \hat{b} = (b_\rho, b_\varphi, b_z)$,

$$|\hat{b}|^2 = b_\rho^2 + b_\varphi^2 + b_z^2 = 1. \tag{1.3}$$

A particular case of the metric (1.1) is the Kerr metric in Kerr-Schield coordinates (cf. [14], [15]):

$$K = \frac{2mr^3}{r^4 + a^2 z^2}, \quad b_\rho = \frac{r\rho}{r^2 + a^2}, \quad b_\varphi = \frac{a\rho}{r^2 + a^2}, \quad b_z = \frac{z}{r}, \tag{1.4}$$

where $r(\rho, z)$ is defined by the relation

$$\frac{\rho^2}{r^2 + a^2} + \frac{z^2}{r^2} = 1. \tag{1.5}$$

Note that the ergosphere (outer ergosphere) for the Kerr metric is

$$K = \frac{2mr^3}{r^4 + a^2 z^2} = 1,$$

and the event horizons (outer and inner event horizons) are

$$r_\pm = m \pm \sqrt{m^2 - a^2}, \quad 0 < a < m. \tag{1.6}$$

Substituting $r = r_\pm$ in (1.5) we see that the equation of the event horizon is an ellipse in (ρ, z) coordinates.

Note that we do not consider the full Kerr spacetime but only a part restricted to $\mathbb{R}^3 \times \mathbb{R}$ (cf. [6], [15]).

Another example of the metric of the form (1.1) is the Gordon's metric (cf. [8], [9]) describing the propagation of light in a moving dielectric medium:

$$dx_0^2 - \sum_{j=1}^{3} dx_j^2 + (n^2 - 1)\Big(\sum_{j=0}^{3} v^{(j)} dx_j\Big)^2, \tag{1.7}$$

where $n(x)$ is the reflection index, $v^{(0)} = \frac{1}{(1-\frac{|w|^2}{c^2})^{\frac{1}{2}}}$, $v^{(j)} = \frac{w_j}{(1-\frac{|w|^2}{c^2})^{\frac{1}{2}}}$, $1 \le j \le 3$, $w(x) = (w_1, w_2, w_3)$ is the velocity of the flow, c is the speed of light in the vacuum.

In this paper we shall pay a special attention to the acoustic metric describing the acoustic waves in a moving fluid flow (cf. [13]). Assuming, for the simplicity of notation, that the speed of the sound and the density are equal to 1, we have the following Hamiltonian in the polar coordinates (ρ, φ)

$$H = \Big(\xi_0 + \frac{A}{\rho}\xi_\rho + \frac{B}{\rho^2}\xi_\varphi\Big)^2 - \xi_\rho^2 - \frac{1}{\rho^2}\xi_\varphi^2, \tag{1.8}$$

where

$$v = \frac{A}{\rho}\hat{\rho} + \frac{B}{\rho}\hat{\varphi} \tag{1.9}$$

is the velocity of the flow, $\hat{\rho} = (\cos\varphi, \sin\varphi)$, $\hat{\varphi} = (-\sin\varphi, \cos\varphi)$, i.e. $\frac{A}{\rho}\hat{\rho}$ is the radial component of v and $\frac{B}{\rho}\hat{\varphi}$ is the angular component of v.

This paper studies the phenomenon of superradiance for the spacetimes having a black hole and ergosphere. The region between the ergosphere and the event horizon is called the ergoregion. We briefly describe the main contributions to the superradiance phenomenon.

On the level of particles this phenomenon was discovered by R. Penrose [11].

He proposed the following thought experiment:

Suppose a particle with an energy ε_0 enters the ergoregion and somehow splits into two particles with energies ε_1 and ε_2. By the conservation of energy $\varepsilon_0 = \varepsilon_1 + \varepsilon_2$. It may happen inside the ergoregion that $\varepsilon_2 < 0$. The particle with the negative energy enters the black hole at some time and the particle with positive energy travels to the infinity. Since $\varepsilon_2 < 0$ we have that $\varepsilon_1 = \varepsilon_0 - \varepsilon_2 > \varepsilon_0$, i.e. the particle leaving the ergoregion has an energy larger than ε_0.

This phenomenon is called the extraction of energy from the black hole.

On the level of the waves, i.e. solutions of the wave equation, Ya. Zeldovich [16] and his then student A. Starobinsky [12] considered the radial component of a single mode of the wave equation after the separation of variables. For this radial component they studied the scattering on the real line and found that the outgoing energy flux may be larger than the incoming energy flux. This is the superradiance on the level of modes. Finster, Kamran, Smoller and Yau [7] gave a far reaching generalization of the result of [12] to the level of wave packets. They constructed a solution of the Cauchy problem for the wave equation with the initial data far from the ergosphere such that the restriction of the energy to the complement of a neighborhood of the black hole has the limit when $x_0 \to +\infty$ larger than the energy of the initial data. Recently, Dafermas, Rodnianski and Shlyapentoch [2] undertook a comprehensive study of the time-dependent scattering theory in the case of the Kerr metric. As a by-product of this theory they obtained the superradiance phenomenon. A comprehensive survey of the results on the superradiance is given in a recent book by Brito, Cardoso and Pani [1].

Our approach to the superradiance phenomenon is in the spirit of the Penrose approach.

In §2 we consider the Cauchy problem for the wave equation with highly oscillatory initial data containing a large parameter k and supported in a small neighborhood U_0 of some point P_0 in the ergoregion. The highly oscillatory initial data allow to construct a geometric optics type solution $u(x_0, x)$ of the Cauchy problem concentrated in a neighborhood of two null-geodesics γ_+ and γ_- starting at the point P_0. In §3 we get that the energy integral $E_{x_0}(u)$ is equal to $E_{x_0}(u_0^+) + E_{x_0}(u_0^-)$ modulo lower order terms in k, where u_0^+ and u_0^- are supported in a neighborhood of γ_+ and γ_-, respectively.

When the neighborhood U_0 is contained in the ergoregion we can arrange the initial data in U_0 such that $E_{x_0}(u_0^-) < 0$ and therefore $E(u_0^+) > E_0(u)$ for large k, i.e. the superradiance takes place. Note that the results of §2 and §3 can be applied to a general wave equation (2.7) corresponding to a general Lorentzian metric having an ergorigion. In §4 we study the acoustic metric with the angular velocity much larger than the radial velocity (see (1.9)). We prove that u_0^+ escapes to the infinity when $x_0 \to +\infty$ and u_0^- crosses the event horizon in a finite time. In §5 we study the acoustic metric when the angular component of the velocity flow is less than the radial component, i.e. $|B| < |A|$. In this case both u_0^+ and u_0^- cross the event horizon in a finite time.

In §6 we consider the case of a white hole, i.e. $A > 0$. Note that the reversal of the time x_0 to $-x_0$ transforms the black hole into a white hole. Therefore, if we consider the Cauchy problem with the same initial data on $(-\infty, 0]$ instead of $[0, +\infty)$ we get that u_0^+ tends to infinity when $x_0 \to -\infty$ and u_0^- approaches the event horizon when $x_0 \to -\infty$.

In §7 we consider the case of the Kerr metric assuming that the initial point P_0 and the two null-geodesics γ_+ and γ_- lie in the equatorial plane. This case is similar to the acoustic case when $|B| < |A|$. As in §5 we get that the superradiance phenomenon is "short-lived" since u_0^+ and u_0^- disappear inside the black hole after a finite time.

In §8 we study the behaviour of u_0^+ and u_0^- inside the event horizon and we prove that in the case of the Kerr metric both u_0^+ and u_0^- end on the singularity ring.

Finally, in §9 we study the Kerr metric when $a^2 = m^2$ (the extremal case) and when $a^2 > m^2$ (the case of naked singularity). We show that the behavior of u_0^+ and u_0^- does not change significantly even when there is no black hole.

2 Geometric optics type solutions of the Cauchy problem

Consider the system of null-bicharacteristics for the Hamiltonian (1.2):

(2.1)

$$\frac{dx_0}{ds} = \frac{\partial H}{\partial \xi_0},\ x_0(0) = 0,\ \frac{d\rho}{ds} = \frac{\partial H}{\partial \xi_\rho},\ \ \rho(0) = \rho',$$

$$\frac{d\varphi}{ds} = \frac{\partial H}{\partial \xi_\varphi},\ \varphi(0) = \varphi',\ \frac{dz}{ds} = \frac{\partial H}{\partial \xi_z},\ z(0) = z',\ \frac{d\xi_\rho}{ds} = -\frac{\partial H}{\partial \rho},\ \xi_\rho(0) = \eta_\rho,$$

$$\frac{d\xi_\varphi}{ds} = -\frac{\partial H}{\partial \varphi}, \xi_\varphi(0) = \eta_\varphi, \frac{d\xi_z}{ds} = -\frac{\partial H}{\partial z}, \xi_z(0) = \eta_z, \frac{d\xi_0}{ds} = -\frac{\partial H}{\partial x_0} = 0, \xi_0(0) = \eta_0.$$

Note that $\frac{\partial H}{\partial x_0} = 0$ since the metric is stationary.

The bicharacteristic (2.1) is called the null-bicharacteristic if for any $s \in \mathbb{R}$

$$\begin{aligned} H(\rho(s), \varphi(s), z(s), \xi_0(s), \xi_\rho(s), \xi_\varphi(s), \xi_z(s)) \\ = \xi_0^2 - \hat{\xi} \cdot \hat{\xi} + K(-\xi_0 + \hat{b} \cdot \hat{\xi})^2 = 0 \end{aligned} \tag{2.2}$$

along the bicharacteristic. Since the metric is stationary, to have (2.2) for all s it is enough to have

$$\eta_0^2 - \hat{\eta} \cdot \hat{\eta} + K(\rho', \varphi', z')(-\eta_0 + \hat{b}(\rho', \varphi', z') \cdot \hat{\eta})^2 = 0, \tag{2.3}$$

where $\hat{\eta} = (\eta_\rho, \frac{\eta_\varphi}{\rho'}, \eta_z)$.

Equation (2.2) is a quadratic equation in ξ_0:

$$(1+K)\xi_0^2 - 2K(\hat{b} \cdot \hat{\xi})\xi_0 + K(\hat{b} \cdot \hat{\xi})^2 - \hat{\xi} \cdot \hat{\xi} = 0. \tag{2.4}$$

It has two distinct real roots $\xi_0^\pm = \lambda^\pm(\rho, \varphi, z, \xi_\rho, \xi_\varphi, \xi_z)$, where

$$\lambda^\pm = \frac{K\hat{b} \cdot \hat{\xi} \pm \sqrt{\Delta_1}}{1+K}, \quad \lambda^- < \lambda^+, \tag{2.5}$$

$$\Delta_1 = (1+K)\hat{\xi} \cdot \hat{\xi} - K(\hat{b} \cdot \hat{\xi})^2. \tag{2.6}$$

We shall call the null-bicharacteristics corresponding to $\xi_0^\pm = \lambda^\pm$ the "plus" ("minus") null-bicharacteristics, respectively. The projection of null-bicharacteristic on (ρ, φ, z)-space is called null-geodesic.

Consider the wave equation corresponding to the metric (1.1):

$$\Box_g u(x_0, x) \overset{def}{=} \sum_{j,k=0}^{3} \frac{1}{\sqrt{-g(x)}} \frac{\partial}{\partial x_j}\Big(\sqrt{-g}\ g^{jk}(x) \frac{\partial u(x_0, x)}{\partial x_k}\Big) = 0, \tag{2.7}$$

where $[g^{jk}]_{j,k=0}^3$ is the inverse to the metric tensor in (1.1), $g^{00} > 0, g(x) = (\det[g^{jk}]_{j,k=0}^3)^{-1}$. We shall construct geometric optic type solutions with initial value supported in U_0 where U_0 is a small neighborhood of some point P_0 in the ergoregion. Such geometric optics solutions are well-known and can be found in many references (see, for example, the book [3], §64).

Let $\chi_0(\rho, \varphi, z)$ be a C_0^∞ function with the support in U_0 and $y_0 = (\rho_0, \varphi_0, z_0)$ be the coordinates of P_0. Let $y' = (\rho', \varphi', z')$ be any point in U_0, and let $\eta = (\eta_\rho, \eta_\varphi, \eta_z)$.

Denote by

$$x = x^\pm(x_0, y, \eta) \tag{2.8}$$

the solution of (2.1) when $\eta_0^\pm = \lambda^\pm(\rho', \varphi', z', \eta_\rho, \eta_\varphi, \eta_z)$ (cf. (2.5)), i.e. $x = x^+(x_0, y, t)$ is a "plus" null-geodesics and $x = x^-(x_0, y, \eta)$ is the "minus" null-geodesics.

Suppose that the Jacobian

$$\frac{Dx^\pm}{Dy}(x_0, y, \eta) \neq 0 \tag{2.9}$$

for $0 \leq x_0 \leq T^\pm,\ y \in U_0,\ \eta$ is fixed. Then by the inverse function theorem $y = y^\pm(x_0, x, \eta)$ and $y^\pm(0, x, \eta) = x$. The points where $\frac{Dx^\pm}{Dy} = 0$ are called focal points. Assuming (2.9) we have following theorem (cf. [3], §64):

Theorem 2.1. *Let k be a large parameter. There exist solutions $u^\pm(x_0, \rho, \varphi, z)$ of (2.7) having the form*

$$u^\pm = u_0^\pm + \frac{1}{k} v^\pm(x_0, \rho, \varphi, z, k), \tag{2.10}$$

where

$$u_0^\pm = e^{ikS^\pm} a_0^\pm(x_0, \rho, \varphi, z, \eta_\rho, \eta_\varphi, \eta_z), \tag{2.11}$$

v^+ and its derivatives are uniformly bounded on $[0, T^\pm]$, eikonals $S^\pm$ satisfy the equations

$$\begin{aligned} &S^\pm_{x_0} - \lambda^\pm(\rho, \varphi, z, S_\rho, S_\varphi, S_z) = 0, \quad 0 \leq x_0 \leq T^\pm, \\ &S^\pm\big|_{x_0=0} = \rho\eta_\rho + \varphi\eta_\varphi + z\eta_z. \end{aligned} \tag{2.12}$$

and

$$a_0^\pm(\rho, \varphi, z, \eta_\rho, \eta_\varphi, \eta_z) = C^\pm(x_0, x, \eta)\chi_0(y^\pm(x_0, x, \eta)), \tag{2.13}$$

where $C^\pm \neq 0$ and bounded, $C^\pm(0, x, \eta) = 1$.

Moreover, the correction term $v^\pm(x_0, \rho, \varphi, z, \eta_\rho, \eta_\varphi, \eta_z)$ can be chosen such that $u^\pm = u_0^\pm + \frac{1}{k}v^\pm$ satisfies the following initial conditions:

$$u^\pm\big|_{x_0=0} = \chi_0(\rho, \varphi, z)e^{ik(\rho\eta_\rho + \varphi\eta_\varphi + z\eta_z)}, \tag{2.14}$$

$$\frac{\partial u^\pm}{\partial x_0}\Big|_{x_0=0} = ik\lambda^\pm(\rho, \varphi, z, \eta_\rho, \eta_\varphi, \eta_z)\chi_0(\rho, \varphi, z)e^{ik(\rho\eta_\rho + \varphi\eta_\varphi + z\eta_z)} \tag{2.15}$$

It follows from (2.13) that the support of $a_0^\pm$ in (x_0,ρ,φ,z) space is contained in the union $W^\pm$ of all "plus" ("minus") null-geodesics starting on the support of $\chi_0(\rho,\varphi,z)$. Therefore $\operatorname{supp} u_0^\pm \subset W^\pm$ and $\operatorname{supp} u^\pm$ is contained in $W^\pm$ modulo $O(\frac{1}{k})$.

Remark 2.1.
We will need to consider also the case when either $\frac{Dx^+}{Dy}(x_0,y_0,\eta)$ or $\frac{Dx^-}{Dy}(x_0,y_0,\eta)$ vanish at a finite number of points $x_0^{+,k}, 1\leq k\leq r_+$, $x_0^{-,k}, 1\leq k\leq r_-$, i.e. when the focal points are present. There is a generalization of geometric optics construction (2.11) belonging to Maslov (cf [10], see also [3], §66), that allows to construct solutions $u^\pm$ depending on a large parameter k, solving initial value problem (2.14), (2.15) for the equation (2.7) on arbitrary large interval $[0,T]$. In the neighborhood of the focal point $u^\pm = u_0^\pm + \frac{1}{k}v^\pm$, where $u_0^\pm$ has a more complicate form than (2.11) (see [10] and [3], §66 for details). However outside of the neighborhoods of focal points $u_0^\pm$ can again be represented in a form (2.11). It follows from the construction that $\operatorname{supp} u^+ \subset W^+$ modulo terms of order $O\big(\frac{1}{k}\big)$ and $\operatorname{supp} u^- \subset W^-$ modulo $O\big(\frac{1}{k}\big)$ as in the case of the absence of focal points.

Let

$$u = u^+ + u^-. \tag{2.16}$$

Then u is an exact solution of (2.7) satisfying the initial conditions

$$u\big|_{x_0=0} = u^+\big|_{x_0=0} + u^-\big|_{x_0=0} = 2\chi_0 e^{ik(\rho\eta_\rho+\varphi\eta_\varphi+z\eta_z)}, \tag{2.17}$$

$$\frac{\partial u}{\partial x_0}\Big|_{x_0=0} = ik(\lambda^+ + \lambda^-)\chi_0 e^{ik(\rho\eta_\rho+\varphi\eta_\varphi+z\eta_z)}. \tag{2.18}$$

3 Conservation of energy integrals

If $u(x_0,x)$ is the solution of (2.7) with some initial conditions and $u(x_0,x)$ has a compact support in x then

$$E_{x_0}(u) = \frac{1}{2}\int\limits_{\mathbb{R}^3}\Big(g^{00}\Big|\frac{\partial u(x_0,x)}{\partial x_0}\Big|^2 - \sum_{j,k=1}^{3} g^{jk}(x)\frac{\partial u}{\partial x_j}\overline{\frac{\partial u}{\partial x_k}}\Big)\sqrt{g}dx \tag{3.1}$$

is independent of x_0.

To show this take the derivative of the right hand side of (3.1) in x_0. Using that $u(x_0, x)$ satisfies (2.7) and integrating by parts we get that $\frac{d}{dx_0}E_{x_0}(u) \equiv 0$, i.e. $E_{x_0}(u) = E_0(u)$.

In particular, $E_{x_0}(u^\pm) = E_0(u^\pm)$. Using the initial conditions (2.14), (2.15) we get

$$E_0(u^\pm) = \frac{k^2}{2}\int\limits_{\mathbb{R}^3}\Big(g^{00}\Big(\frac{\partial S^\pm}{\partial x_0}\Big)^2 - \sum_{j,k=1}^{3} g^{jk}\frac{\partial S^\pm}{\partial x_j}\frac{\partial S^\pm}{\partial x_k}\Big)\chi_0^2\sqrt{g}dx. \tag{3.2}$$

Note that when $x_0 = 0$ $S^\pm_{x_j} = \eta_j, 1 \leq j \leq 3,\ S^\pm_{x_0} = \lambda^\pm(\rho, \varphi, z, \eta_\rho, \eta_\varphi, \eta_z)$.

Theorem 3.1. *Let the metric $\sum_{j,k=0}^{3} g_{jk}(x)dx_jdx_k$ has a nonempty ergoregion Ω, i.e. the domain where $g_{00}(x) < 0$. Let $P_0 \in \Omega$, $y_0 = (\rho_0, \varphi_0, z_0)$ be the coordinates of P_0, and U_0 is a small neighborhood of P_0 containing in Ω. One always can find $\eta = (\eta_\rho, \eta_\varphi, \eta_z)$ such that*

$$\lambda^+(y_0, \eta) > \lambda^-(y_0, \eta) > 0.$$

Then, assuming that U_0 is small, we have

$$E_{x_0}(u^-) < 0, \quad E_{x_0}(u^+) > 0,$$

and

$$E_{x_0}(u^+) = E_{x_0}(u) - E_{x_0}(u^-) > E_{x_0}(u),$$

where $u = u^+ + u^-$. Thus the superradiance takes place.

Proof: It follows from (2.12) that

$$g^{00}(S^\pm_{x_0})^2 + 2\sum_{j=1}^{3} g^{0j}S^\pm_{x_j}S^\pm_{x_0} + \sum_{j,k=1}^{3} g^{jk}S^\pm_{x_j}S^\pm_{x_k} = 0, \tag{3.3}$$

since $H = (\xi_0 - \lambda^+)(\xi_0 - \lambda^-)$.

Therefore

$$-\sum_{j,k=1}^{3} g^{jk}S^\pm_{x_j}S^\pm_{x_k} = g^{00}(S^\pm_{x_0})^2 + 2\sum_{j=1}^{3} g^{0j}S^\pm_{x_j}S^\pm_{x_0}. \tag{3.4}$$

Substituting (3.4) into (3.2) we get

$$E_0(u^\pm) = k^2\int\limits_{\mathbb{R}^3} S^\pm_{x_0}\Big(g^{00}S^\pm_{x_0} + \sum_{j=1}^{3} g^{0j}\frac{\partial S^\pm}{\partial x_j}\Big)\chi_0^2\sqrt{g}dx. \tag{3.5}$$

It follows from (3.3), analogously to (2.5), (2.6), that

$$g^{00}S^{\pm}_{x_0} + \sum_{j=1}^{3} g^{0j} S^{\pm}_{x_j} = \pm\sqrt{\Delta_1}, \tag{3.6}$$

where

$$\Delta_1 = \Big(\sum_{j=1}^{3} g^{0j} S^{\pm}_{x_j}\Big)^2 - g^{00} \sum_{j,k=1}^{3} g^{jk} S^{\pm}_{x_j} S^{\pm}_{x_k}. \tag{3.7}$$

Note that in the case of metric (1.1) Δ_1 has the form (2.6) and $g^{00} = 1 + K$.

Therefore considering, for definiteness, the case of metric (1.1) we have

$$E_0(u^+) = k^2 \int\limits_{\mathbb{R}^3} S^{+}_{x_0} \sqrt{\Delta_1}\chi_0^2 \sqrt{g}dx = k^2 \int\limits_{\mathbb{R}^3} \lambda^+ \sqrt{\Delta_1}\chi_0^2 \rho d\rho d\varphi dz, \tag{3.8}$$

$$E_0(u^-) = -k^2 \int\limits_{\mathbb{R}^3} S^{-}_{x_0} \sqrt{\Delta_1}\chi_0^2 \sqrt{g}dx = -k^2 \int\limits_{\mathbb{R}^3} \lambda^- \sqrt{\Delta_1}\chi_0^2 \rho d\rho d\varphi dz, \tag{3.9}$$

where $x_0 = 0$ and $S^{\pm}_{x_0}, \Delta^{\pm}_1$ have the form (2.5), (2.6). Note that $E_{x_0}(u^{\pm}) = E_0(u^{\pm})$.

Now compute $E_0(u^+ + u^-)$.

It follows from (2.17), (2.18) and (3.1) that (cf. (2.4))

$$E_0(u^+ + u^-) = \frac{k^2}{2}\int\limits_{\mathbb{R}^3} \Big((1+K)(\lambda^+ + \lambda^-)^2\chi_0^2 - (K(\hat{b}\cdot\hat{\eta})^2 - \hat{\eta}\cdot\hat{\eta})4\chi_0^2\Big)\rho d\rho d\varphi dz. \tag{3.10}$$

Since

$$\lambda^+ + \lambda^- = \frac{2K(\hat{b}\cdot\hat{\eta})}{1+K}, \tag{3.11}$$

we have

(3.12)

$$
\begin{aligned}
E_0(u^+ + u^-) =& \frac{k^2}{2}\int\limits_{\mathbb{R}^3}\Big[(1+K)\frac{4K^2(\hat{b}\cdot\hat{\eta})^2}{(1+K)^2} - 4(K(\hat{b}\cdot\hat{\eta})^2 - \hat{\eta}\cdot\hat{\eta})\Big]\chi_0^2\rho d\rho d\varphi dz \\
=& 2k^2\int\limits_{\mathbb{R}^3}\Big(\frac{K^2(\hat{b}\cdot\hat{\eta})^2}{1+K} - K(\hat{b}\cdot\hat{\eta})^2 + \hat{\eta}\cdot\hat{\eta}\Big)\chi_0^2\rho d\rho d\varphi dz \\
=& 2k^2\int\limits_{\mathbb{R}^3}\frac{(1+K)\hat{\eta}\cdot\hat{\eta} - K(\hat{b}\cdot\hat{\eta})^2}{1+K}\chi_0^2\rho d\rho d\varphi dz \\
=& 2k^2\int\limits_{\mathbb{R}^3}\frac{\Delta_1}{1+K}\chi_0^2\rho d\rho d\varphi dz
\end{aligned}
$$

From the other side,

$$
E_0(u^+) + E_0(u^-) = k^2\int\limits_{\mathbb{R}^3}(\lambda^+\sqrt{\Delta_1} - \lambda^-\sqrt{\Delta_1})\chi_0^2\rho d\rho d\varphi dz. \tag{3.13}
$$

Since $\lambda^+ - \lambda^- = \frac{2\sqrt{\Delta_1}}{1+K}$ we are getting that

$$
E_0(u^+ + u^-) = E_0(u^+) + E_0(u^-),
$$

and thus for any x_0

$$
E_{x_0}(u^+ + u^-) = E_{x_0}(u^+) + E_{x_0}(u^-). \tag{3.14}
$$

When $y = (\rho', \varphi', z')$ is outside of the ergoregion then $K < 1$ and

$$
K(\hat{b}\cdot\hat{\eta})^2 - \hat{\eta}\cdot\hat{\eta} < 0
$$

for all $\hat{\eta} \neq 0$. Therefore λ_+ and λ_- have opposite signs (cf. (2.4)).

When $y_0 = (\rho_0, \varphi_0, z_0)$ belongs to the ergoregion then $K > 1$ and one can choose $\hat{\eta}_0 = (\eta_\rho, \eta_\varphi, \eta_z)$ such that

$$
K(\hat{b}\cdot\hat{\eta}_0)^2 - \hat{\eta}_0\cdot\hat{\eta}_0 > 0.
$$

Thus $\lambda^+(y_0, \hat{\eta}_0)$ and $\lambda^-(y_0, \hat{\eta}_0)$ have the same sign.

Without loss of generality we can assume that $\lambda_+(y_0, \hat{\eta}_0) > \lambda^-(y_0, \hat{\eta}_0) > 0$. Otherwise we replace $\hat{\eta}_0$ by $-\hat{\eta}_0$ and use that $\lambda^+(y_0, \hat{\eta}_0) = -\lambda^-(y_0, -\hat{\eta}_0)$.

Then $E_{x_0}(u^-) < 0$ when $\operatorname{supp}\chi_0$ contains in a small neighborhood of (ρ_0, φ_0, z_0) (see (3.9)). ☐

Since $E_{x_0}(u^+ + u^-) = E_{x_0}(u^+) + E_{x_0}(u^-)$ we have that

$$E_{x_0}(u^+) = E_{x_0}(u^+ + u^-) - E_{x_0}(u^-) > E_{x_0}(u^+ + u^-) = E_0(u^+ + u^-). \tag{3.15}$$

Note that $\operatorname{supp} u_0^{\pm} \subset W^{\pm}$ (cf. (2.13)), where $W^+(W^-)$ is the union of all null-geodesics starting at (y, η) where $y \in \operatorname{supp}\chi_0$. Therefore, for any x_0 one can find k so large that

$$E_{x_0}(u_0^+) > E_{x_0}(u^+ + u^-) = E_0(u^+ + u^-). \tag{3.16}$$

We took into account (3.15) and that $\frac{E_{x_0}(u^+)}{E_{x_0}(u_0^+)} = 1 + O\big(\frac{1}{k}\big)$. Therefore (3.16) show that the superradiance takes place. Note that the "minus" null-geodesics can not leave the ergoregion since $\xi_0^- = \lambda^-(y, \eta) > 0$. It will be shown that when $x_0 > T_1$ u_0^- is inside the event horizon. Therefore, outside the event horizon $\frac{E_{x_0}(u^-)}{E_{x_0}(u_0^-)} = O\big(\frac{1}{k}\big)$ for x_0 large.

Remark 3.1. The case of real-valued solutions. Since coefficients of equation (2.7) are real, the real (and imaginary) part of complex-valued solution also satisfies (2.7). Let $u^{\pm} = u_0^{\mp} + \frac{1}{k}v^{\pm}$ be the same as in (2.10). Denote by $u_R^{\pm} = u_{0R}^{\mp} + \frac{1}{k}v_R^{\pm}$ the real part of (2.10). Since $a_0^{\pm}$ is real-valued we have

$$u_R^{\pm} = \frac{1}{2}(u^{\pm} + \overline{u^{\pm}}) \equiv a_0^{\pm} \cos kS^{\pm} + O\Big(\frac{1}{k}\Big). \tag{3.17}$$

Moreover, (2.14), (2.15) imply

(3.18)

$$\begin{aligned}
&u_R^{\pm}\big|_{x_0=0} = \chi_0(\rho, \varphi, z) \cos k(\rho\eta_\rho + \varphi\eta_\varphi + z\eta_z),\\
&\frac{\partial u_R^{\pm}}{\partial x_0}\Big|_{x_0=0} = -k\lambda^{\pm}(\rho, \varphi, z, \eta_r, \eta_\varphi, \eta_z)\chi_0(\rho, \varphi, z) \sin k(\rho\eta_\rho + \varphi\eta_\varphi + z\eta_z).
\end{aligned}$$

Since (3.1) holds for real-valued solutions too and since (3.1) is independent of x_0, we have, as in (3.2):

(3.19) $E_0(u_R^\pm)$

$$= \frac{k^2}{2} \int\limits_{\mathbb{R}^3} \Big(g^{00} \Big(\frac{\partial S^\pm}{\partial x_0} \Big)^2 - \sum_{j,k=1}^{3} g^{jk} \frac{\partial S^\pm}{\partial x_j} \frac{\partial S^\pm}{\partial x_k} \Big) \chi_0^2 \sin^2 k(\rho\eta_\rho + \varphi\eta_\varphi + z\eta_z)\rho d\rho d\varphi.$$

Therefore, as in (3.8), (3.9), we get

$$E_0(u_R^+) = k^2 \int\limits_{\mathbb{R}^3} \lambda^+ \sqrt{\Delta_1} \sin^2 k(\rho\eta_\rho + \varphi\eta_\varphi + z\eta_z)\chi_0^2 \rho d\rho d\varphi dz > 0, \tag{3.20}$$

$$E_0(u_R^-) = -k^2 \int\limits_{\mathbb{R}^3} \lambda^- \sqrt{\Delta_1} \sin^2 k(\rho\eta_\rho + \varphi\eta_\varphi + z\eta_z)\chi_0^2 \rho d\rho d\varphi dz < 0. \tag{3.21}$$

Since $\sin^2 k(\rho\eta_\rho + \varphi\eta_\varphi + z\eta_z) = \frac{1}{2} - \frac{1}{2}\cos 2k(\rho\eta_\rho + \varphi\eta_\varphi + z\eta_z)$ and since integrals containing $\cos 2k(\rho\eta_\rho + \varphi\eta_\varphi + z\eta_z)$ are decaying fast when $k \to \infty$ we have, as in (3.14), (3.15), (3.16), that

$$E_{x_0}(u_{0R}^+) > E_{x_0}(u_R^+ + u_R^-) = E_0(u_R^+ + u_R^-) \tag{3.22}$$

when k is large.

Thus superradiance holds also for real-valued solutions. Note that

$$E_{x_0}(u) = E_{x_0}(u_R) + E_{x_0}(u_I),$$

where $u = u_R + iu_I, , u_R = \Re u, u_I = \Im u$. Also $E_{x_0}(u_R) \approx \frac{1}{2}E_{x_0}(u),\ E_{x_0}(u_I) \approx \frac{1}{2}E_{x_0}(u)$ when $k \to \infty$.

4 Superradiance in the case of the acoustic metric

Consider in more details the case of the acoustic metric (1.8) (cf. [13], [4], [5]). The equation $H = 0$ (cf. (1.8)) has two roots

$$\xi_0^\pm = -\frac{A}{\rho}\xi_\rho - \frac{B}{\rho^2}\xi_\varphi \pm \sqrt{\Delta_1}, \tag{4.1}$$

where

$$\Delta_1 = \xi_\rho^2 + \frac{1}{\rho^2}\xi_\varphi^2. \tag{4.2}$$

The black hole is $\{\rho < |A|,\ A < 0\}$, and the ergoregion is $\{|A| < \rho < \sqrt{A^2 + B^2}\}$. We assume, for definiteness, $B > 0$.

We shall find first simple expressions for the differential equations for the "plus" and "minus" null-geodesics. Note that

$$\begin{aligned} H_{\xi_\rho} &= 2\Big(\xi_0 + \frac{A}{\rho}\xi_\rho + \frac{B}{\rho^2}\xi_\varphi\Big)\frac{A}{\rho} - 2\xi_\rho, \\ H_{\xi_0} &= 2\Big(\xi_0 + \frac{A}{\rho}\xi_\rho + \frac{B}{\rho^2}\xi_\varphi\Big). \end{aligned}$$

It follows from (2.1) that in the acoustic case

$$\frac{d\rho^\pm}{dx_0} = \frac{H_{\xi_\rho}}{H_{\xi_0}} = \frac{A}{\rho} - \frac{\xi_\rho}{\xi_0^\pm + \frac{A}{\rho}\xi_\rho + \frac{B\xi_\varphi}{\rho^2}} = \frac{(\frac{A^2}{\rho^2} - 1)\xi_\rho + \frac{A}{\rho}(\xi_0^\pm + \frac{B\xi_\varphi}{\rho^2})}{\xi_0^\pm + \frac{A}{\rho}\xi_\rho + \frac{B\xi_\varphi}{\rho^2}}. \tag{4.3}$$

Since $H = 0$, the following quadratic equation for ξ_p holds:

$$\Big(\frac{A^2}{\rho^2} - 1\Big)\xi_\rho^2 + 2\frac{A}{\rho}\Big(\xi_0^\pm + \frac{B\xi_\varphi}{\rho^2}\Big)\xi_\rho + \Big(\xi_0^\pm + \frac{B\xi_\varphi}{\rho^2}\Big)^2 - \frac{1}{\rho^2}\xi_\varphi^2 = 0. \tag{4.4}$$

Therefore

$$\xi_\rho = \frac{-\frac{A}{\rho}\big(\xi_0^\pm + \frac{B\xi_\varphi}{\rho^2}\big) \pm \sqrt{\Delta_2^\pm}}{\frac{A^2}{\rho^2} - 1}, \tag{4.5}$$

where

$$\begin{aligned} \Delta_2^\pm &= \frac{A^2}{\rho^2}\Big(\xi_0^\pm + \frac{B\xi_\varphi}{\rho^2}\Big)^2 - \Big(\frac{A^2}{\rho^2} - 1\Big)\Big(\xi_0^\pm + \frac{B\xi_\varphi}{\rho^2}\Big)^2 + \Big(\frac{A^2}{\rho^2} - 1\Big)\frac{\xi_\varphi^2}{\rho^2} \\ &= \Big(\xi_0^\pm + \frac{B\xi_\varphi}{\rho^2}\Big)^2 + \Big(\frac{A^2}{\rho^2} - 1\Big)\frac{\xi_\varphi^2}{\rho^2}. \end{aligned} \tag{4.6}$$

Substituting (4.5) in the numerator of (4.3) we get

$$\frac{d\rho^\pm}{dx_0} = \frac{\pm\sqrt{\Delta_2^\pm}}{\xi_0^\pm + \frac{A}{\rho}\xi_r + \frac{B\xi_\varphi}{\rho^2}} = \frac{\pm\sqrt{\Delta_2^\pm}}{\pm\sqrt{\xi_\rho^2 + \frac{1}{\rho^2}\xi_\varphi^2}}. \tag{4.7}$$

Therefore

$$\frac{d\rho^+}{dx_0} = \frac{\pm\sqrt{\Delta_2^\pm}}{\sqrt{\Delta_1}}, \qquad \frac{d\rho^-}{dx_0} = \frac{\pm\sqrt{\Delta_2^\pm}}{-\sqrt{\Delta_1}} \tag{4.8}$$

In this section we consider the case when the angular velocity is much larger than the radial velocity, i.e. $B \gg |A|$. The following theorem holds:

Theorem 4.1. *Let* $2|A| < \rho_0 < \sqrt{A^2+B^2}$. *Choose* $\eta_\rho = -\frac{2|A|}{\rho_0}, \eta_\varphi = -\rho_0\sqrt{1-\frac{4A^2}{\rho_0^2}}$, *i.e.* $\eta_\rho^2 + \frac{\eta_\varphi^2}{\rho_0^2} = \frac{4A^2}{\rho_0^2} + \left(1-\frac{4A^2}{\rho^2}\right) = 1$. *Suppose*

$$B > \left(1+\frac{2A^2}{\rho_0^2}\right)\frac{\rho_0}{\sqrt{1-\frac{4A^2}{\rho_0^2}}}. \tag{4.9}$$

Let $u^\pm = u_0^\pm + \frac{1}{k}v^\pm$ *be the same as in (2.11), (2.12) satisfying the initial conditions (2.14), (2.15).*

Then the superradiance takes place, i.e. $E_{x_0}(u_0^+) > E_0(u^+ + u^-)$ *for large* k, *supp* u_0^+ *tends to infinity when* $x_0 \to +\infty$, *supp* u_0^- *crosses the event horizon* $\rho = |A|$ *when* x_0 *increases.*

Proof: Find first the sign of $\frac{d\rho^\pm}{dx_0}$ at the initial point (ρ_0, φ_0), $\xi_\rho = \eta_\rho$, $\xi_\varphi = \eta_\varphi$ when $x_0 = 0$. Since $\rho_0 > 2|A|$, $\eta_\rho = -\frac{2|A|}{\rho_0}$, $\eta_\varphi = -\rho_0\sqrt{1-\frac{4A^2}{\rho_0^2}}$, we have

$$\xi_0^-\Big|_{x_0=0} = -\frac{A}{\rho_0}\eta_\rho - \frac{B}{\rho_0^2}\eta_\varphi - \sqrt{\Delta_1} = -\frac{2A^2}{\rho_0^2} + \frac{B}{\rho_0}\sqrt{1-\frac{4A^2}{\rho_0^2}} - 1. \tag{4.10}$$

If (4.9) holds then $\xi_0^- > 0$ and therefore the superradiance will take place. Also

$$\frac{d\rho^+}{dx_0}\Big|_{x_0=0} = \frac{A}{\rho_0} - \frac{\eta_\rho}{\sqrt{\eta_\rho^2 + \frac{\eta_\varphi^2}{\rho_0^2}}} = -\frac{|A|}{\rho_0} + 2\frac{|A|}{\rho_0} > 0, \tag{4.11}$$

since $\sqrt{\eta_\rho^2 + \frac{\eta_\varphi^2}{\rho_0^2}} = 1$, $\eta_\rho = -\frac{2|A|}{\rho_0}$. Therefore

$$\frac{d\rho^+}{dx_0} = \frac{\sqrt{\Delta_2^+}}{\sqrt{\Delta_1}} > 0 \ \text{ for all } \ x_0 \geq 0, \tag{4.12}$$

if we can prove that $\Delta_2^+ > 0$ for all $\rho \geq \rho_0$, (zeros of $\Delta_2^\pm$ are called the turning points).

Rewrite equation $\Delta_2^\pm = 0$ (cf. (4.6)) in the form

$$\frac{(A^2+B^2)\xi_\varphi^2}{\rho^4} + \frac{2\xi_0^\pm B\xi_\varphi - \xi_\varphi^2}{\rho^2} + (\xi_0^\pm)^2 = 0, \tag{4.13}$$

where

$$\xi_0^\pm\big|_{x_0=0} = -\frac{2A^2}{\rho_0^2} - \frac{B\eta_\varphi}{\rho_0^2} \pm 1. \tag{4.14}$$

Solving the quadratic equation in $\frac{1}{\rho^2}$ we get

$$\frac{1}{\rho^2} = \frac{-2\xi_0^\pm B\xi_\varphi + \xi_\varphi^2 \pm \sqrt{\delta}}{2(A^2+B^2)\xi_\varphi^2} \tag{4.15}$$

where

$$\begin{aligned} \delta &= (2B\xi_\varphi\xi_0^\pm - \xi_\varphi^2)^2 - 4(A^2+B^2)\xi_\varphi^2(\xi_0^\pm)^2 \\ &= -4B\xi_\varphi^3\xi_0^\pm + \xi_\varphi^4 - 4A^2\xi_\varphi^2(\xi_0^\pm)^2. \end{aligned} \tag{4.16}$$

Since B is large (cf. (4.10), $\xi_\varphi = \eta_\varphi < 0$ and $\xi_0^\pm$ has the form (4.14), we have

$$\delta = 4\xi_\varphi^2|B\xi_\varphi|\xi_0^\pm - 4A^2\xi_\varphi^2(\xi_0^\pm)^2 + \xi_\varphi^4 = 4\xi_\varphi^2\frac{(B\xi_\varphi)^2}{\rho_0^2} - 4\xi_\varphi^2\frac{A^2(B\xi_\varphi)^2}{\rho_0^4} + \cdots,$$

where $\cdots$ means nonsignificant terms. Therefore

$$\sqrt{\delta} = 2\frac{B\xi_\varphi^2}{\rho_0}\Big(1 - \frac{A^2}{\rho_0^2}\Big)^{\frac{1}{2}} + \cdots \tag{4.17}$$

Substituting in (4.15) and taking into the account that $B \gg |A|$ we obtain

$$\begin{aligned} \frac{1}{\rho^2} =& \frac{2|B\xi_\varphi|\big(\frac{B|\xi_\varphi|}{\rho_0^2} - \frac{2A^2}{\rho_0^2} \pm 1\big) \pm \frac{2B\xi_\varphi^2}{\rho_0}\big(1 - \frac{A^2}{\rho_0^2}\big)^{\frac{1}{2}}}{2B^2\xi_\varphi^2} + \cdots \\ =& \frac{1}{\rho_0^2} + \frac{1}{B|\xi_\varphi|}\Big(\pm 1 - \frac{2A^2}{\rho_0^2}\Big) \pm \frac{\big(1 - \frac{A^2}{\rho_0^2}\big)^{\frac{1}{2}}}{B\rho_0} + \cdots \end{aligned} \tag{4.18}$$

Note that $|\xi_\varphi| = \rho_0\big(1 - \frac{4A^2}{\rho_0^2}\big)^{\frac{1}{2}}$.

It follows from (4.18) that $\Delta_2^+ = 0$ has turning points $\rho_2^+ < \rho_1^+ < \rho_0$, where ρ_j^+ solves

$$\frac{1}{(\rho_j^+)^2} = \frac{1}{\rho_0^2} + \frac{1}{B|\xi_\varphi|}\Big(1 - \frac{2A^2}{\rho_0^2}\Big) \pm \frac{\big(1 - \frac{A^2}{\rho_0^2}\big)^{\frac{1}{2}}}{B\rho_0} + \dots, \quad j = 1, 2, \tag{4.19}$$

ρ_1^+ corresponding to the minus sign in (4.19) and ρ_2^+ to the plus sign. Analogously, $\Delta_2^- = 0$ has turning points $\rho_0 < \rho_1^- < \rho_2^-$, where ρ_j^- solves

$$\frac{1}{(\rho_j^-)^2} = \frac{1}{\rho_0^2} + \frac{1}{B|\xi_\varphi|}\Big(- 1 - \frac{2A^2}{\rho_0^2}\Big) \pm \frac{\big(1 - \frac{A^2}{\rho_0^2}\big)^{\frac{1}{2}}}{B\rho_0} + \dots, \quad j = 1, 2. \tag{4.20}$$

Therefore $\Delta_2^+ > 0$ for all $\rho > \rho_1^+$ and hence $\frac{d\rho^+}{dx_0} > 0$ for all $x_0 > 0$. Thus $\rho^+(x_0) \to +\infty$ when $x_0 \to +\infty$.

Note that $\frac{d\rho^-}{dx_0}\big|_{x_0=0} = -\frac{|A|}{\rho_0} + \frac{\eta_\rho}{\sqrt{\eta_\rho^2 + \frac{\eta_\varphi^2}{\rho_0^2}}} = -\frac{3|A|}{\rho_0} < 0$.

Thus $\Delta_2^- > 0$ for $\rho < \rho_1^-$. We have that

$$\frac{d\rho^-}{dx_0} = -\frac{\sqrt{\Delta_2^-}}{\sqrt{\Delta_1}} < 0 \quad \text{for} \quad x_0 > 0, \tag{4.21}$$

and $\rho^-(x_0)$ crosses the event horizon when x_0 increases. Let

$$u^\pm = e^{ikS^\pm} a_0^\pm + \frac{1}{k} v^\pm(x_0, \rho, \varphi, k) \tag{4.22}$$

be the solutions (2.10), (2.11) in the case of the acoustic metric and let $E_{x_0}(u^\pm)$ be the corresponding energy integrals (cf. (3.8), (3.9)).

When the condition (4.9) holds then $\rho^+(x_0) \to +\infty$ if $x_0 \to +\infty$. Hence $\operatorname{supp} u_0^+$ tends to the infinity when $x_0 \to +\infty$. Also $\operatorname{supp} u_0^-$ enters the black hole $\rho = |A|$ when x_0 increases. As in (3.16) we have $E_{x_0}(u_0^+) > E_0(u^+ + u^-)$ for large k, i.e. the superradiance takes place. □

Remark 4.1. The behavior of $\mathbf{u_0^+}$ and $\mathbf{u_0^-}$ on the time interval $(-\infty, \mathbf{0})$.

Consider the behavior of $\rho^+(x_0)$ and $\rho^-(x_0)$ on the time interval $(-\infty, 0]$.

Since $\rho_1^+ < \rho_0$ is the turning point where $\Delta_2^+ = 0$, we get that $\rho^+(x_0)$ decreases when x_0 decreases until it reaches ρ_1^+ at some time $x_0^{(1)} < 0$. Note that the null-geodesics γ^+ has a singularity at $\rho^+(x_0^{(1)}) = \rho_1^+$ since $\sqrt{\Delta_2^+} = C\sqrt{\rho^+(x_0) - \rho_1^+}$ for $x_0 > x_0^{(1)}$. Passing the point $x_0^{(1)}$ we have $\frac{d\rho^+(x_0)}{dx_0} = -\frac{\sqrt{\Delta_2^+}}{\sqrt{\Delta_1^+}} < 0$ for $x_0 < x_0^{(1)}$ (cf. [4]), i.e. $\rho^+(x_0)$ turns and starts increasing when $x_0 < x_0^{(1)}$ decreasing. Since $\Delta_2^+ > 0$ on $(\rho_1^+, +\infty)$ we get that $\rho^+(x_0) \to +\infty$ when $x_0 \to -\infty, x_0 < x_1^{(0)}$. Similarly, for $\rho^-(x_0)$ the point $\rho_1^- > \rho_0$ is also the turning point and $\rho^-(x_0)$ increases when x_0 decreases until it reaches ρ_1^- at some time $x_0^{(2)} < 0$. Then $\rho^-(x_0)$ turns and starts to decrease when x_0 decreases. When $x_0 \to -\infty \quad \rho^-(x_0)$ tends to the event horizon $\rho = |A|$ and $\rho^-(x_0)$ spirals around the event horizon as $x_0 \to -\infty$ (cf. [4]).

Turning points are singularities (focal) points for null-geodesics. When the null-geodesics is passing through the focal point one needs the Maslov extension of the geometric optics construction as stated in Remark 2.1. Note that the turning points ρ_1^+ and ρ_1^- are the simplest type of singularities called the fold singularities. A detailed geometric optics construction for the case of such singularities is presented in [3], Example 66.1.

Since $\rho^+(x_0) \to \infty$ when $x_0 \to -\infty$ we have that $\operatorname{supp} u_0^+$ tends to ∞ when $x_0 \to -\infty$. Also $\operatorname{supp} u_0^-$ tends to the event horizon when $x_0 \to -\infty$ since $\rho^-(x_0)$ approaches $\rho = |A|$ when $x_0 \to -\infty$.

Remark 4.2. The case of acoustic metric with $\mathbf{A = 0}$ (the case of naked singulirity).
Consider the acoustic metric (1.8) when $A = 0$. We assume as above $B > 0, \eta_\varphi < 0$. The equation of the ergosphere is $\rho = B$ and there is no event horizon, i.e. $\rho = 0$ is a "naked" singularity.

We choose the initial point ρ_0 such that $\xi_0^- = -\frac{B\eta_\varphi}{\rho_0^2} - \sqrt{\eta_\rho^2 + \frac{\eta_\varphi^2}{\rho_0^2}} > 0$. Thus

$$\frac{(B^2 - \rho_0^2)\eta_\varphi^2}{\rho_0^4} > \eta_\rho^2. \tag{4.23}$$

It follows from (4.23) that $\rho_0 < B$, i.e ρ_0 belongs to the ergoregion.

We choose $\eta_\rho < 0$ small and η_ρ satisfies (4.21) and choose η_φ such that

$$\frac{\eta_\varphi^2}{\rho_0^2} + \eta_\rho^2 = 1. \tag{4.24}$$

Then

$$\xi_0^{\pm} = \frac{B|\eta_\varphi|}{\rho_0^2} \pm 1.$$

We shall show that the superradiance takes place when $A = 0$ despite the absence of the black hole.

Theorem 4.2. *Suppose* $A = 0, B > 0, \rho_0, \eta_\varphi, \eta_\rho$ *are such that* $\rho_0 < B, \eta_\rho < 0, \eta_\varphi < 0$ *and (4.23), (4.24) hold. Let* $u_0^{\pm} = u_0^{\pm} + \frac{1}{k}v^{\pm}$ *be the same as in Theorem 4.1 with* $\rho_0, \eta_\rho, \eta_\varphi$ *chosen as above and* φ_0 *is arbitrary. Then* $E_0(u^-) < 0$ *since* $\xi_0^- = -\frac{B\eta_\varphi}{\rho_0^2} - \sqrt{\eta_\rho^2 + \frac{\eta_\varphi^2}{\rho_0^2}} > 0$*. Thus* $E_{x_0}(u_0^+) > E_0(u^+ + u^-)$ *for large* k*, i.e. the superradiance holds. Moreover,* u_0^+ *escapes to the infinity when* $x_0 \to +\infty$ *and* u_0^- *approach the singularity* $\rho = 0$ *when* x_0 *increases.*

Proof: The equations (4.3) have the form

$$\frac{d\rho^{\pm}(x_0)}{dx_0} = \frac{-\xi_\rho}{\xi_0^{\pm} + \frac{B\xi_\varphi}{\rho^2}} = -\frac{\xi_\rho}{\pm\sqrt{\Delta_1}}, \tag{4.25}$$

where $\Delta_1 = \xi_\rho^2 + \frac{\xi_\varphi^2}{\rho^2}$. As in (4.5), (4.6), (4.8), we have

$$\Delta_2^{\pm} = \left(\xi_0^{\pm} + \frac{B\xi_\varphi}{\rho^2}\right)^2 - \frac{\xi_\varphi^2}{\rho^2}, \quad \xi_\rho^{\pm} = \mp\sqrt{\Delta_2^+}, \tag{4.26}$$

$$\frac{d\rho^+}{dx_0} = \frac{\mp\sqrt{\Delta_2^+}}{\sqrt{\Delta_1}}, \qquad \frac{d\rho^-}{dx_0} = \frac{\mp\sqrt{\Delta_2^-}}{-\sqrt{\Delta_1}}. \tag{4.27}$$

The turning points, i.e. the solutions of $\Delta_2^{\pm} = 0$ are the solutions of the equations (4.15), (4.16) with $A = 0$. Therefore (cf. (4.18))

$$\frac{1}{\rho^2} = \frac{1}{\rho_0^2} \pm \frac{1}{B|\eta_\varphi|} \pm \frac{1}{B\rho_0} + \dots$$

Note that $|\eta_\varphi| = \rho_0(1 - \eta_\rho^2)^{\frac{1}{2}} < \rho_0$.

Hence, as in (4.21), (4.22), $\Delta_2^+ = 0$ for $\rho_2^+ < \rho_1^+ < \rho_0$ and $\Delta_2^- = 0$ for $\rho_0 < \rho_1^- < \rho_2^-$. At the point $x_0 = 0$ we have $\frac{d\rho^+}{dx_0} = -\eta_\rho > 0$ and $\frac{d\rho^-}{dx_0} = \frac{-\eta_\rho}{-1} < 0$. Therefore $\frac{d\rho^+}{dx_0} = \frac{\sqrt{\Delta_2^+}}{\sqrt{\Delta_1}}$ for $x_0 > 0$, $\frac{d\rho^-}{dx_0} = -\frac{\sqrt{\Delta_2^-}}{\sqrt{\Delta_1}}$ for $x_0 > 0$.

Since $\Delta_2^+ > 0$ on $[\rho_0, +\infty)$ we have that $\rho^+(x_0) \to \infty$ when $x_0 \to +\infty$. Also since $\Delta_2^- > 0$ for $\rho < \rho_0$ we have that $\rho^-(x_0) \to 0$ when x_0 increases.

It follows from (4.26) that $\Delta_2^- \sim \frac{B^2\xi_\varphi^2}{\rho^4}$, $\Delta_1 \sim \frac{B^2\xi_\varphi^2}{\rho^4}$. Therefore $\frac{d\rho^-}{dx_0} \sim -1$ when $\rho \to 0$, i.e. $\rho^-(x_0)$ reaches 0 in a finite time when x_0 increases. Note that

$$\frac{d\varphi}{d\rho} = \frac{-\frac{\xi_\varphi}{\rho^2} + (\xi_0^- + \frac{B\xi_\varphi}{\rho^2})\frac{B}{\rho^2}}{\sqrt{\Delta_2^-}} \sim \frac{\frac{B^2}{\rho^4}\xi_\varphi}{\frac{|B\xi_\varphi|}{\rho^2}} \sim \frac{-B}{\rho^2}.$$

Hence $\varphi \to \infty$ when $\rho \to 0$. Therefore γ_- approaches $\rho = 0$ spiraling when $\rho \to 0$.

Let $u^\pm = u_0^\pm + \frac{1}{k}v^\pm$ be the same as in Theorem 4.1 with $\rho_0, \eta_\varphi, \eta_\rho$ as in (4.23), (4.24). Since $\xi_0^- = -\frac{B\eta_\varphi}{\rho_0^2} - \sqrt{\eta_\rho^2 + \frac{\eta_\varphi^2}{\rho_0^2}} > 0$ we have that $E_0(u^-) < 0$ and $E_{x_0}(u_0^+) > E_0(u^+ + u^-)$ for large k as in §3, i.e. the superradiance take place. Since $\rho^+(x_0) \to \infty$ when $x_0 \to +\infty$ we have that u_0^+ escapes to the infinity when $x_0 \to +\infty$. Also since $\rho^-(x_0) \to 0$ spiraling when x_0 increases, we have that u_0^- approaches the singularity $\rho = 0$ when x_0 increases. $\square$

Remark 4.3. Behavior of $\mathbf{u_0^+}$ and $\mathbf{u_0^-}$ on the time interval $(-\infty, \mathbf{0}]$ in the case of naked singularity.
As in Remark 4.1 we can study first the behavior of $\rho^+(x_0)$ and $\rho^-(x_0)$ when $x_0 \in (-\infty, 0]$. We get as in Remark 4.1 that $\rho^+(x_0)$ decays on $[x_0^{(1)}, 0]$ where $x_0^{(1)} < 0$. $x_0^{(1)}$ is the turning point, i.e. $\rho^+(x_0^{(1)}) = \rho_1^+$. Then $\rho^+(x_0)$ increasing for $x_0 < x_0^{(0)}$ and tends to $+\infty$ when $x_0 \to -\infty$. Analogously, $\rho^-(x_0)$ increases on $[x_0^{(2)}, 0]$ where $\rho^-(x_0^{(2)}) = \rho_1^-$ is the turning point and then $\rho^-(x_0)$ decreases for $x_0 < x_0^{(2)}$ when x_0 decreases. Note that the equation of $\rho^-(x_0)$ for $x_0 < x_0^{(0)}$ is $\frac{d\rho^-}{dx_0} = \frac{\sqrt{\Delta_2^-}}{\sqrt{\Delta_1}}$. Therefore $\frac{d\rho^-}{dx_0} \sim +1$ when $\rho \to 0$ and $\rho^-(x_0)$ reaches 0 in a finite time when x_0 decreases. Also $\frac{d\varphi}{d\rho} = O\left(\frac{1}{\rho^2}\right)$. Thus $\varphi \to \infty$ when $\rho \to 0$, i.e. $\rho^-(x_0)$ is approaching 0 spiraling.

Since the null-geodesics γ_+ and γ_- are passing through the turning points ρ_1^+ and ρ_1^-, respectively, we need the Remark 2.1 to construct the solutions u_0^+ and u_0^- near turning points. Since $\rho^+(x_0) \to +\infty$ when $x_0 \to -\infty$ we get that u_0^+ escapes to the infinity when $x_0 \to -\infty$. Also u_0^- approaches the singularity $\rho = 0$ when x_0 decreases since $\rho^-(x_0) \to 0$ when x_0 decreases.

5 The acoustic metric when $|B| \leq |A|$

Let $|A| < \rho_0 < \sqrt{A^2 + B^2}$, $A < 0, B > 0$, i.e. (ρ_0, φ) belongs to the ergoregion. Suppose

$$B < |A|. \tag{5.1}$$

If $(\eta_\rho, \eta_\varphi)$ are such that

$$\lambda^- = -\frac{A\eta_\rho}{\rho_0} - \frac{B\eta_\varphi}{\rho_0^2} - \sqrt{\eta_\rho^2 + \frac{\eta_\varphi^2}{\rho_0^2}} > 0, \tag{5.2}$$

then $\eta_\rho > 0$. Indeed,

$$\frac{|B\eta_\varphi|}{\rho_0^2} \leq \frac{B}{|A|}\frac{|\eta_\varphi|}{\rho_0} < \sqrt{\eta_\rho^2 + \frac{\eta_\varphi^2}{\rho_0^2}},$$

and η_ρ must be positive.

Theorem 5.1. *Suppose* $0 < B \leq |A|$. *Choose* $\eta_\rho = \frac{|A|}{\rho_0}, \eta_\varphi = -B$. *Let* $u^\pm = u_0^\pm + \frac{1}{k}v^\pm$ *be the same as in Theorem 4.1 with the choice of* $|A| < \rho_0 < \sqrt{A^2 + B^2}, \eta_\rho = \frac{|A|}{\rho_0}, \eta_\varphi = -B$. *Then* $E_0(u_0^-) < 0, E_{x_0}(u_0^+) > E_0(u^+ + u^-)$ *for large* k, *i.e. the superradiance takes place. Moreover* u_0^+ *and* u_0^- *cross the event horizon in a finite time. Therefore the superradiance is "short-lived" outside the event horizon.*

Proof: We have

$$\lambda^- = \frac{A^2}{\rho_0^2} + \frac{B^2}{\rho_0^2} - \sqrt{\frac{A^2}{\rho^2} + \frac{B^2}{\rho_0^2}} > 0 \tag{5.3}$$

since $\rho_0^2 < A^2 + B^2$. It follows from (5.3) that the superradiance takes place.

As in (4.8) we have

$$\frac{d\rho^+}{dx_0} = \frac{\pm\sqrt{\Delta_2^+}}{\sqrt{\Delta_1}}, \quad \frac{d\rho^-}{dx_0} = \frac{\pm\sqrt{\Delta_2^-}}{-\sqrt{\Delta_1}},$$

where $\rho^\pm(x_0)$ are the same as in (4.3), Δ_1 and $\Delta_2^\pm$ are the same as in (4.2), (4.6).

We shall show that if (5.1) holds then $\Delta_2^+ > 0$ for all $\rho > |A|$, i.e. $\rho^+(x_0)$ has no turning points. We get from (4.16), with $\xi_\varphi = -B$ and $\xi_0^\pm$ as in (5.3) that

$$\delta = 4B^4\xi_0^+ + B^4 - 4A^2B^2(\xi_0^+)^2 = B^2(4B^2\xi_0^+ + B^2 - 4A^2(\xi_0^+)^2).$$

The roots of the quadratic equation $B^2+4B^2t-4A^2t^2 = 0$ are $t = \frac{2B^2\pm\sqrt{4B^4+4A^2B^2}}{4A^2}$. The largest root $t_1 = \frac{B^2+\sqrt{B^4+A^2B^2}}{2A^2} < \frac{A^2+\sqrt{A^4+A^2A^2}}{2A^2} = \frac{1+\sqrt{2}}{2}$. From other side $\xi_0^+ = \frac{A^2+B^2}{\rho_0^2} + \sqrt{\frac{A^2+B^2}{\rho_0^2}} > 1 + 1 = 2 > \frac{1+\sqrt{2}}{2}$. Thus $\xi_0^+ > t_1$ and consequently $\delta < 0$. Therefore $\Delta_2^+ > 0$ for all ρ, i.e. $\rho^+(x_0)$ has no turning points. Consider $\frac{d\rho^+}{dx_0}$ when $x_0 = 0$. We have

$$\frac{d\rho^+}{dx_0}\Big|_{x_0=0} = -\frac{|A|}{\rho_0} - \frac{\frac{|A|}{\rho_0}}{\sqrt{\frac{A^2}{\rho_0^2} + \frac{B^2}{\rho_0^2}}} < 0. \tag{5.4}$$

Therefore $\rho^+(x_0)$ decreases until it reaches the event horizon. The solution $\rho^-(x_0)$ can not leave the ergoregion since $\lambda^- < 0$ outside of the ergoregion and it always hits the event horizon in finite time. In the all cases of the acoustic metric $\rho^-(x_0)$ increases from $-\infty$ to some point $x_0^{(2)}$ where it reaches the turning point ρ_- inside the ergoregion. For $x_0 > x_0^{(2)}$ it decreases until it reaches the event horizon (cf. [4]). Thus u_0^+ and u_0^- also reach the event horizon in finite time.

Therefore there is a superradiance when $B < |A|$ but it is "short-lived" since both u_0^+ and u_0^- disappear inside the black hole after a finite time.

6 The superradiance in the case of white hole

Consider the acoustic metric (1.8) when $A > 0$. Then $\rho = A$ will be a white hole (cf [13]). Note that the change in the sign of B has no effect whether there is a black or white hole. From the other side the change of the sign of both A and B is equivalent to the change of the sign of ξ_0 in (1.8), i.e. to the reversal of the time from x_0 to $-x_0$. Therefore the study of the superradiance for the case of the white hole is equivalent to the study of the superradiance for the black hole on the time interval $(-\infty, 0]$.

Theorem 6.1. *Consider the acoustic metric (1.8) with $A > 0$. Then $\rho = A$ is a white hole. Let $u^\pm = u^\pm + \frac{1}{k}v^\pm$ are the same as in (2.10)–(2.15). When* *the point* (ρ_0, φ_0) *is inside the ergoregion,* $A < \rho_0 < \sqrt{A^2 + B^2}$ *one can choose* $(\eta_\rho, \eta_\varphi)$ *such that the superradiance takes place (cf. §3),* $\operatorname{supp} u_0^+$ *tends to the infinity when* $x_0 \to +\infty$ *and* $\operatorname{supp} u_0^-$ *tends to the event horizon spiraling when* $x_0 \to +\infty$*.*

Proof: Instead of giving a direct proof of Theorem 6.1 we change x_0 to $-x_0$, i.e. we consider the case of black hole on $(-\infty, 0]$.

It was shown in §5 that $\rho^+(x_0)$ with the initial data $\rho^+(0) = \rho_0$ has no turning points. Therefore since (5.4) holds and $\Delta_2^+ > 0$ we have $\rho^+(x_0) \to +\infty$ when $x_0 \to -\infty$. Also in the case when (4.10) holds we have that $\rho^+(x_0)$ with the initial data $\rho^+(0) = \rho_0$ has a turning point $\rho_1^+ < \rho_0$. Therefore $\rho^+(x_0)$ decreases when x_0 decreases until it reaches the turning point $\rho = \rho_1^+$. After this $\rho^+(x_0) \to +\infty$ when $x_0 \to -\infty$. Therefore in both cases when either (4.10) or (5.1) are satisfied $\rho^+(x_0) \to +\infty$ when $x_0 \to -\infty$. Thus the support of corresponding u_0^+ tends to the infinity when $x_0 \to -\infty$. In both §4 and §5 $\rho^-(x_0)$ approaches the event horizon spiraling when $x_0 \to -\infty$ and therefore $\operatorname{supp} u_0^-$ also approaches the event horizon.

Changing back x_0 to $-x_0$ we prove Theorem 6.1.

7 The case of the Kerr metric

In this section we consider the superradiance for the case of the Kerr metric (1.2), (1.4). Note that $b_z = \frac{z}{r} = 0$ when $z = 0$. Therefore we have from (2.1) that $z \equiv 0$ implies

$$\frac{dz}{ds} = H_{\xi_z} = -2\xi_z + 2K(-\xi_0 + \hat{b} \cdot \hat{\xi})b_z \equiv 0, \tag{7.1}$$

and therefore $\xi_z \equiv 0$.

Therefore if the initial point $P_o = (\rho_0, \varphi_0, 0)$ is in the equatorial plane $z = 0$ and if $\xi_z = 0$ then the bicharacteristic system (2.1) is reduced to the bicharacteristic system for the Hamiltonian

$$H(\rho, \varphi, 0, \xi_\rho, \xi_\varphi, 0) = \xi_0^2 - \xi_\rho^2 - \frac{1}{\rho^2}\xi_\varphi^2 + K\Big(- \xi_0 + b_\rho \xi_\rho + b_\varphi \frac{\xi_\varphi}{\rho}\Big)^2 = 0, \tag{7.2}$$

and the null-geodesics lie in the plane $z = 0$.

As in (2.5), (2.6)

$$\lambda^\pm(y_0, \eta') = \frac{Kb' \cdot \eta' \pm \sqrt{\Delta_1}}{1 + K} \tag{7.3}$$

and

$$\Delta_1 = (1+K)\xi'\cdot\xi' - K(b'\cdot\xi')^2,$$

where $z=0,\ b'=(b_\rho, b_\varphi),\ \eta'=(\eta_\rho, \frac{\eta_\varphi}{\rho}),\ y_0=(\rho_0,\varphi_0,0)$.

We will take point $y_0=(\rho_0,\varphi_0,0)$ in the ergoregion and choose $\eta'=(\eta_\rho, \frac{1}{\rho_0}\eta_\varphi)$ such that

$$\lambda^-(y_0,\eta')>0.$$

It follows from (1.4) for $z=0$

$$b_\rho = \frac{\sqrt{\rho^2-a^2}}{\rho},\quad b_\varphi=\frac{a}{\rho},\quad K=\frac{2m}{\sqrt{\rho^2-a^2}}, \tag{7.4}$$

since $\rho^2=r^2+a^2$ when $z=0$.

We choose

$$\eta_\rho = b_\rho(y_0),\quad \eta_\varphi=\rho_0 b_\varphi(y_0),\quad z=0. \tag{7.5}$$

Note that

$$\eta_\varphi = \rho_0\frac{a}{\rho_0}=a. \tag{7.6}$$

Since

$$b'(y_0)\cdot\eta' = b_\rho^2+b_\varphi^2=1, \tag{7.7}$$

we have

$$\Delta_1 = (1+K)(\eta'\cdot\eta') - K(b'\cdot\eta')^2 = (1+K)-K=1. \tag{7.8}$$

Therefore

$$\lambda^\pm(y_0,\eta') = \frac{K\pm 1}{K+1}>0, \tag{7.9}$$

since $K>1$ in the ergoregion. Thus $\lambda^+=1,\ \lambda^-=\frac{K-1}{K+1}<1$.

Theorem 7.1. *Consider the Kerr metric (1.1). Let* $u^\pm = u_0^\pm + \frac{1}{k}v^\pm$ *be the same as in (2.10)–(2.15). Choose initial point* $y_0=(\rho_0,\varphi_0,0)$ *in the ergoregion. Choose* $\eta_\rho=b_\rho(y_0), \eta_\varphi=\rho_0 b_\varphi(y_0), \eta_z=0, z=0$ *(cf. (7.4)). Thus* $\lambda^-(y_0,\eta_\rho,\eta_\varphi)=\frac{K-1}{K+1}>0$. *Therefore the superradiance takes place. Moreover* u_0^+ *and* u_0^- *disappear inside the outer event horizon in a finite time, i.e. the superradiance is "short-lived".*

We shall study first the null-bicharacteristic (2.1) corresponding to the root $\xi_0^+ = \lambda^+(y_0, \eta')$, starting at (y_0, η'). We have

$$\frac{d\rho^+}{ds} = \frac{\partial H}{\partial \xi_\rho} = -2\xi_\rho + 2K(-\xi_0^+ + b'\cdot\xi')b_\rho, \quad \rho^+(0) = \rho_0, \tag{7.10}$$

$$\frac{dx_0}{ds} = \frac{\partial H}{\partial \xi_0} = 2\xi_0^+ - 2K(-\xi_0^+ + b'\cdot\xi'), \quad x_0(0) = 0, \tag{7.11}$$

$$\frac{d\varphi}{ds} = H_{\xi_\varphi} = -\frac{2\xi_\varphi}{\rho^2} + 2K(-\xi_0^+ + b'\cdot\xi')\frac{b_\varphi}{\rho}. \tag{7.12}$$

We have (see (2.5))

$$\frac{\partial H}{\partial \xi_0} = 2(1+K)\xi_0^+ - 2Kb'\cdot\xi' = 2\sqrt{\Delta_1}, \tag{7.13}$$

where Δ_1 is the same as in (2.6) for $z = 0, \xi_z = 0$.

Also we have

$$\frac{\partial H}{\partial \xi_\rho} = -2\xi_\rho + 2Kb_\rho(-\xi_0^+ + b'\cdot\xi') = 2(Kb_\rho^2 - 1)\xi_\rho + 2Kb_\rho\Big(-\xi_0^+ + b_\varphi\frac{\xi_\varphi}{\rho}\Big). \tag{7.14}$$

Write (7.2) as the quadratic equation for ξ_ρ:

$$(Kb_\rho^2 - 1)\xi_\rho^2 + 2K\Big(-\xi_0^+ + b_\varphi\frac{\xi_\varphi}{\rho}\Big)b_\rho\xi_\rho + K\Big(-\xi_0^+ + b_\varphi\frac{\xi_\varphi}{\rho}\Big)^2 + (\xi_0^+)^2 - \frac{\xi_\varphi^2}{\rho^2} = 0. \tag{7.15}$$

Therefore

$$\xi_\rho = \frac{-Kb_\rho\Big(-\xi_0^+ + b_\varphi\frac{\xi_\varphi}{\rho}\Big) \pm \sqrt{\Delta_2^+}}{Kb_\rho^2 - 1} \tag{7.16}$$

where

$$\begin{aligned} \Delta_2^+ =& K^2b_\rho^2\Big(-\xi_0^+ + b_\varphi\frac{\xi_\varphi}{\rho}\Big)^2 - (Kb_\rho^2 - 1)\Big[K\Big(-\xi_0^+ + b_\varphi\frac{\xi_\varphi}{\rho}\Big)^2 + (\xi_0^+)^2 - \frac{\xi_\varphi^2}{\rho^2}\Big] \\ =& K\Big(-\xi_0^+ + b_\varphi\frac{\xi_\varphi}{\rho}\Big)^2 - (Kb_\rho^2 - 1)\Big((\xi_0^+)^2 - \frac{\xi_\varphi^2}{\rho^2}\Big). \end{aligned} \tag{7.17}$$

It follows from (7.9), that (7.17) simplifies to

$$\Delta_2^+ = \frac{\rho^2 - a^2}{\rho^2}. \tag{7.18}$$

Substituting (7.16) into (7.14) we get

$$\frac{\partial H}{\partial \xi_\rho} = \pm 2\sqrt{\Delta_2^+}. \tag{7.19}$$

Therefore

$$\frac{d\rho^+}{dx_0} = \frac{H_{\xi_\rho}}{H_{\xi_0}} = \frac{\pm\sqrt{\Delta_2^+}}{\sqrt{\Delta_1^+}}. \tag{7.20}$$

Thus $\rho = \rho^+(x_0)$ is either increasing or decreasing depending on the sign of $\frac{\partial H}{\partial \xi_\rho}$.

It is enough to check the sign of $\frac{\partial H}{\partial \xi_\rho}$ at the initial point $x_0 = 0$.

Having η_ρ, η_φ as in (7.5) we get

$$\frac{\partial H}{\partial \xi_\rho} = -2b_\rho(y_0) + 2Kb_\rho\Big(-\lambda^+ + b_\rho\eta_\rho + b_\varphi\frac{\eta_\varphi}{\rho_0}\Big). \tag{7.21}$$

It follows from (7.21)

$$\frac{\partial H}{\partial \xi_\rho}(y_0, \eta) = -2b_\rho(y_0) < 0,$$

since $-\lambda^+ + b_\rho^2 + b_\varphi^2 = 0$.

Therefore, $\frac{d\rho^+}{dx_0} < 0$ for all x_0 and $\rho = \rho^+(x_0)$ will decay when x_0 increases and will approach the outer event horizon.

Consider now the "minus" null-geodesics starting at (y_0, η'). We have as before $\frac{d\rho^-}{dx_0} = \pm\frac{\sqrt{\Delta_2^-}}{\sqrt{\Delta_1^-}}$ where Δ_2^- is the same as Δ_2^+ with $\xi_0^+ = 1$ replaced by $\xi_0^- = \frac{K-1}{K+1}$.

To determine the sign of $\frac{d\rho^-}{dx_0}$ consider the initial point $y_0 = (\rho_0, \varphi_0, 0)$, $\eta' = (b_\rho, \rho_0 b_\varphi)$. We have $\frac{\partial H}{\partial \xi_0} = -2\sqrt{\Delta_1}$ and

$$\frac{\partial H}{\partial \xi_\rho} = -2\eta_\rho + 2Kb_\rho(-\lambda^- + b'\cdot\eta') = -2b_\rho + \frac{4Kb_\rho}{1+K} > 0. \tag{7.22}$$

Since $\frac{\partial H}{\partial \xi_0} < 0, \frac{\partial H}{\partial \xi_\rho} > 0$ we get that

$$\frac{d\rho^-}{dx_0} = -\frac{\sqrt{\Delta_2^-}}{\sqrt{\Delta_1}} < 0. \tag{7.23}$$

Therefore $\rho^-(x_0)$ also approaches the event horizon when x_0 increases.

Consider the behavior of $\rho^+(x_0)$ and $\rho^-(x_0)$ when they approach the outer event horizon $r = r_+$ where $r = \sqrt{\rho^2 - a^2}$, $r_\pm = m \pm \sqrt{m^2 - a^2}$. Note that

$$\begin{aligned} 1 - Kb_\rho^2 &= 1 - \frac{2m}{r}\Big(\frac{r}{\sqrt{r^2+a^2}}\Big)^2 \\ &= \frac{r^2 + a^2 - 2mr}{r^2+a^2} = \frac{(r-r_+)(r-r_-)}{r^2+a^2}, \end{aligned}$$

i.e. $1 - Kb_\rho^2 = 0$ when $r = r_+$. We have $\Delta_2^+ \sim K\big(-1 + \frac{a^2}{\rho^2}\big)^2$ when $r \to r_+$. Note that (7.16) can be rewritten in the form $H_{\xi_\rho} = \pm 2\sqrt{\Delta_2}$.

Since $H_{\xi_\rho} < 0$ for $\rho^+(x_0)$ at $x_0 = 0$ (cf. (7.21)) we take $-\sqrt{\Delta_2^+}$ in (7.16). Therefore when $\rho \to \hat{\rho}_+ = \sqrt{a^2 + r_+^2}$ we have canceling of $\sqrt{K}b_\rho - 1$:

$$\begin{aligned} (7.24)\quad \xi_\rho^+ &= \frac{-Kb_\rho\big(-1 + \frac{a^2}{\rho^2}\big) - \sqrt{\Delta_2^+}}{Kb_\rho^2 - 1} \\ &\sim \frac{Kb_\rho\big(1 - \frac{a^2}{\rho^2}\big) - \sqrt{K}\big(1 - \frac{a^2}{\rho^2}\big)}{(\sqrt{K}b_\rho - 1)(\sqrt{K}b_\rho + 1)} = \frac{\sqrt{K}\big(1 - \frac{a^2}{\rho^2}\big)}{\sqrt{K}b_\rho + 1}. \end{aligned}$$

Therefore ξ_ρ^+ has a finite nonzero limit when $r \to r_+$ and therefore $\frac{d\rho^+}{dx_0}$ has nonzero limit when $r \to r_+$. Hence $\rho^+(x_0)$ crosses the event horizon when x_0 increases. Also $H_\rho > 0$ for $\rho^-(x_0)$ at $x_0 = 0$ (see (7.22)). Therefore we have $+\sqrt{\Delta_2^-}$ in (7.16) for $\rho^-(x_0)$. Note that $\xi_0^- = \frac{K-1}{K+1} = \frac{2m - r_0}{2m + r_0}$ where $r_0 = \sqrt{\rho_0^2 - a^2} > r_+$. Since $\frac{2m - r_0}{2m + r_0} < \frac{2m - r_+}{2m + r_+}$ when $r_+ < r_0 < 2m$ and $\frac{2m - r_+}{2m + r_+} \le \frac{a^2}{a^2 + r_+^2}$ we get that $-\xi_0^- + \frac{a^2}{\rho^2} > 0$ when $\rho = \sqrt{a^2 + r_+^2}$. When $\rho \to \sqrt{a^2 + r_+^2}$ we have

(7.25)

$$\xi_\rho^- = \frac{-Kb_\rho\big(-\xi_0^- + \frac{a^2}{\rho^2}\big) + \sqrt{\Delta_2^-}}{Kb_\rho^2 - 1} \sim \frac{-Kb_\rho\big(-\xi_0^- + \frac{a^2}{\rho^2}\big) + \sqrt{K}\big(-\xi_0^- + \frac{a^2}{\rho^2}\big)}{Kb_\rho^2 - 1}.$$

Canceling $-\sqrt{K}b_\rho + 1$ we get that ξ_ρ^- has a finite nonzero limit when $r \to r_+$. Therefore as in the case of $\rho^+(x_0)$ we have that $\rho^-(x_0)$ crosses the outer event horizon when x_0 increases. Since $\rho^+(x_0)$ and $\rho^-(x_0)$ cross the event horizon in a finite time we get that u_0^+ and u_0^- also cross the outer event horizon in a finite time. Thus the superradiance for the Kerr metric is short-lived as in

§5. $\square$

Remark 7.1. The behavior of $\mathbf{u_0^+}$ and $\mathbf{u_0^-}$ on $(-\infty, \mathbf{0}]$.
Consider the behavior of $\rho^-(x_0)$ for $x_0 < 0$. Note that $\rho^-(x_0)$ has a turning point $\rho_*, \rho_0 < \rho_* < \sqrt{a^2 + 4m^2}, \Delta_2^-(\rho_*) = 0$ because $\Delta_2^-(\rho_0) > 0$ (cf. (7.22)) and $\Delta_2^- < 0$ when $\hat{\rho} = \sqrt{4m^2 + a^2}$ where $\rho = \hat{\rho}$ is the ergosphere. Indeed, $K = 1$ on the ergosphere and

(7.26)

$$\Delta_2^- = \Big(-\xi_0^- + \frac{a^2}{\hat{\rho}^2}\Big)^2 + (1 - b_\rho^2)\Big((\xi_0^-)^2 - \frac{a^2}{\hat{\rho}^2}\Big) = (\xi_0^-)^2 - 2\xi_0^- \frac{a^2}{\hat{\rho}^2} + (\xi_0^-)^2 \frac{a^2}{\rho^2}.$$

Thus

$$\Delta_2^- = \xi_0^- \Big(1 + \frac{a^2}{\hat{\rho}^2}\Big)\Big[\xi_0^- - \frac{2a^2}{4m^2 + a^2}\Big].$$

We have

$$\xi_0^- = \frac{2m - r_0}{2m + r_0} < \frac{2m - r_+}{2m + r_+} = \frac{m - \sqrt{m^2 - a^2}}{3m + \sqrt{m^2 - a^2}} \le \frac{a^2}{4m^2 - a^2} < \frac{2a^2}{4m^2 + a^2}$$

since $a < m$. Hence $\Delta_2^- < 0$ on the ergosphere.

Note that $\rho^-(x_0)$ increases when x_0 decreases from 0 to $x_0^{(1)}$, i.e. $\rho^-(x_0^{(1)})$ is the turning point. Then the sign of $\sqrt{\Delta_2^-}$ changes, i.e. we change $+\sqrt{\Delta_+^-}$ to $-\sqrt{\Delta_+^-}$ in (7.16) and the equation for $\rho^-(x_0)$ becomes

$$\frac{d\rho^-}{dx_0} = \frac{\sqrt{\Delta_2^-}}{\sqrt{\Delta_1^-}} \quad \text{for} \quad x_0 < x_0^{(1)}. \tag{7.27}$$

When $r \to r_+$ we get $\xi_\rho^- \sim \frac{-2K(-\xi_0^- + \frac{a^2}{\rho^2})}{Kb_\rho^2 - 1} \sim \frac{1}{r - r_+}$. Therefore

$$\Delta_1^- = (1 + K)\Big((\xi_\rho^-)^2 + \frac{a^2}{\rho^2}\Big) - K\Big(b_\rho \xi_\rho^- + \frac{a^2}{\rho^2}\Big) \sim \frac{1}{(r - r_+)^2} \tag{7.28}$$

when $r \to r_+$, or $\rho \to \hat{\rho}_+ = \sqrt{r_+^2 + a^2}$. Therefore

$$\frac{d\rho^-}{dx_0} \le C|\rho - \hat{\rho}_+|. \tag{7.29}$$

Hence $\ln|\rho - \hat{\rho}_+| \le Cx_0 + C_1$, and

$$0 < \rho - \hat{\rho}_+ \le C_2 e^{Cx_0}.$$

Thus $\rho \to \hat{\rho}_+$ when $x_0 \to -\infty$. It follows from (7.12) that

$$\frac{d\varphi}{d\rho} = O\Big(\frac{1}{r - r_+}\Big). \tag{7.30}$$

Therefore $\varphi \to \infty$ when $\rho \to \hat{\rho}_+$. Thus the null-geodesics γ_- approaches the event horizon $\rho = \hat{\rho}_+$ spiraling when $x_0 \to -\infty$. Note that $\rho^+(x_0) \to \infty$ when $x_0 \to -\infty$ since $\Delta_2^+ = \frac{\rho^2 - a^2}{\rho^2} > 0$ for all $\rho > a$ and $\frac{d\rho^+}{dx_0} = -\frac{\sqrt{\Delta_2^+}}{\sqrt{\Delta_1^+}} < 0$.

Remark 7.2.
The case of the white Kerr hole arises when the time direction is reversed (cf. §6). It is equivalent to studying the black Kerr hole on $(-\infty, 0]$. This study was done in Remark 7.1.

Thus we have that $\operatorname{supp} u_0^+$ tends to infinity when $x_0 \to -\infty$. Since γ_- spirals approaching $\rho = \hat{\rho}_+$ when $x_0 \to -\infty$, we have that $\operatorname{supp} u_0^-$ also approaches $\rho = \hat{\rho}_+$ when $x_0 \to -\infty$.

8 The behavior of solutions inside the Kerr outer horizon

8.1 The case of null-geodesics in the equatorial plane

Consider what happens with the "plus" and "minus" null-geodesics γ_+ and γ_- after they cross the outer event horizon $r_+ = m + \sqrt{m^2 - a^2}$.

Theorem 8.1. *After crossing the outer event horizon both null-geodesics γ^+ and γ^- cross also the inner event horizon and end on the singularity ring $\rho = a$ at a finite time (see Fig. 1).*

Consider first the case of the "plus" null-geodesics. We have, as in §7:

$$\frac{d\rho^+(x_0)}{dx_0} = -\frac{\sqrt{\Delta_2^+}}{\sqrt{\Delta_1^+}}, \tag{8.1}$$


γ_+
P_0
γ_-
$\rho = a$
$\rho = \sqrt{a^2 + r_-^2}$
$\rho = \sqrt{a^2 + r_+^2}$
$\rho = \sqrt{a^2 + 4m^2}$

Figure 1: The case $a < m$.
Null-geodesics γ_+ starts at P_0 and reaches $\rho = a$ at a finite time $x_0^{(1)} > 0$. When $x_0 \to -\infty$ γ_+ tends to infinity. Null-geodesics γ_- starts at P_0 and reaches $\rho = a$ at finite time $x_0^{(2)} > 0$. When $x_0 < 0$ γ_- passes a turning point and then spirals, approaching the outer event horizon.

where $\Delta_2^+ = \frac{\rho^2-a^2}{\rho^2}$,

$$\Delta_1^+ = (1+K)\Big((\xi_\rho^+)^2 + \frac{\xi_\varphi^2}{\rho^2}\Big) - K\Big(b_\rho\xi_\rho^+ + b_\varphi\frac{\xi_\varphi}{\rho}\Big)^2. \tag{8.2}$$

Since $\Delta_2^+ \neq 0, \Delta_1^+ \neq 0$ we get from (8.1) that $\rho^+(x_0)$ decreases when x_0 increases. After γ_+ cross the outer event horizon it crosses also the inner event horizon since (7.24) holds near $r = r_0$. Then it is approaching the ring singularity $(\rho - a)^2 + z^2 = 0$

We shall study the behavior of γ_+ near the ring singularity, i.e. when $z = 0,\ \rho$ is close to $a,\ \rho > a$. It follows from (7.16) that

$$\xi_\rho^+ = \frac{-\frac{2m}{\sqrt{\rho^2-a^2}}\frac{\sqrt{\rho^2-a^2}}{\rho}\big(-1+\frac{a^2}{\rho^2}\big) + \frac{\sqrt{\rho^2-a^2}}{\rho}}{\frac{2m}{\sqrt{\rho^2-a^2}}\big(\frac{\sqrt{\rho^2-a^2}}{\rho}\big)^2 - 1}. \tag{8.3}$$

Hence

$$\begin{aligned}\Delta_1^+ = \Big(1+\frac{2m}{\sqrt{\rho^2-a^2}}\Big)(1+O(\rho^2-a^2)) - \frac{2m}{\sqrt{\rho^2-a^2}}(1+O(\rho^2-a^2))\\ = 1 + O(\sqrt{\rho^2-a^2}).\end{aligned} \tag{8.4}$$

Thus $\frac{d\rho^+}{dx_0} = -\frac{\sqrt{\Delta_2^+}}{\sqrt{\Delta_1^+}} \leq -C\sqrt{\rho-a}$. Therefore

$$\frac{d\rho}{\sqrt{\rho-a}} \leq -Cdx_0. \tag{8.5}$$

Integrating (8.5) from x_0 to t, where $x_0 < t,\ \rho^+(t) - a > 0$, we get

$$2\sqrt{\rho^+(t)-t} - 2\sqrt{\rho^+(x_0)-a} \leq -C(t-x_0). \tag{8.6}$$

Increasing t further we get t_0 such that $\rho^+(t_0) - a = 0$, i.e. $\rho^+(x_0)$ reaches $\rho = a$ at $x_0 = t_0$.

Note that (cf. (7.12)

$$\frac{d\varphi}{d\rho} = \frac{H_{\xi_\varphi}}{H_{\xi_\rho}} = \frac{-\frac{2\xi_\varphi}{\rho^2} + 2K\big(-\xi_0^+ + b_\rho\xi_\rho^+ + b_\varphi\frac{\xi_\varphi}{\rho}\big)\frac{b_\varphi}{\rho}}{\pm\sqrt{\Delta_2^+}} = \frac{C(\rho)}{(\rho^2-a^2)^{\frac{1}{2}}}, \tag{8.7}$$

where $C(\rho) \neq 0$. Hence $\varphi(x_0) \to \varphi_0$ when $\rho \to a$. Therefore the "plus" null-geodesics γ_+ ends at the point (a, φ_0) of the singularity ring at some time t_0.

Now consider the case of "minus" null-geodesics γ_-. It also crosses the inner event horizon when x_0 increases since (7.25) holds. Note that $0 < \xi_0^- = \lambda^- < 1$ (cf. (7.9)). We have (cf. (7.17))

$$
\begin{aligned}
(8.8) \qquad \Delta_2^- &= K\Big(- \xi_0^- + \frac{\rho^2}{a^2}\Big)^2 - (Kb_\rho^2 - 1)\Big((\xi_0^-)^2 - \frac{a^2}{\rho^2}\Big) \\
&= \frac{2m}{\sqrt{\rho^2 - a^2}}\Big((-\xi_0^- + 1) + \frac{\rho^2 - a^2}{a^2}\Big)^2 \\
&\quad - \Big(\frac{2m}{\sqrt{\rho^2 - a^2}}\Big(\frac{\sqrt{\rho^2 - a^2}}{\rho}\Big)^2 - 1\Big)\Big((\xi_0^-)^2 - 1 + \frac{\rho^2 - a^2}{a^2}\Big)^2 \\
&= \frac{2m(1 - \xi_0^-)^2}{\sqrt{\rho^2 - a^2}}(1 + O(\sqrt{\rho^2 - a^2})).
\end{aligned}
$$

Also we have (cf. (7.16)

$$
(8.9) \quad \xi_\rho^- \quad = \quad \frac{-Kb_\rho\big(- \xi_0^- + \frac{a^2}{\rho^2}\big) + \sqrt{\Delta_2^+}}{Kb_\rho^2 - 1} \quad = \quad +\sqrt{\Delta_2^-} \ + \ O(1).
$$

Hence (cf. (8.2)

(8.10)

$$
\begin{aligned}
\Delta_1^- &= \Big(1 + \frac{2m}{\sqrt{\rho^2 - a^2}}\Big)\Big(\Big(\xi_\rho^-\Big)^2 + \frac{a^2}{\rho^2}\Big) - \frac{2m}{\sqrt{\rho^2 - a^2}}\Big(\frac{\sqrt{\rho^2 - a^2}}{\rho}\xi_\rho^- + \frac{a^2}{\rho^2}\Big)^2 \\
&= \frac{(2m)^2(1 - \xi_0^-)^2}{\rho^2 - a^2} + O\Big(\frac{1}{\sqrt{\rho^2 - a^2}}\Big).
\end{aligned}
$$

It follows from (7.22) and (8.8), (8.10) that

$$
(8.11) \qquad \frac{d\rho^-}{dx_0} = -\frac{\sqrt{\Delta_2^-}}{\sqrt{\Delta_1^-}} \leq -\frac{C_1(\rho^2 - a^2)^{-\frac{1}{4}}}{C_2(\rho^2 - a^2)^{-\frac{1}{2}}} \leq -C_4(\rho - a)^{\frac{1}{4}},
$$

i.e.

$$
(\rho - a)^{-\frac{1}{4}} d\rho \leq -C_4 dx_0.
$$

Analogously to (8.6), integrating from x_0 to t_1, $t_1 > x_0$, $\rho^-(t_1) - a > 0$, we get that there exists $t^{(0)}$ such that $\rho^-(t^{(0)}) - a = 0$, i.e. γ_- also ends on $\rho = a$.

8.2 Behavior of $\mathbf{u}_0^+$ and $\mathbf{u}_0^-$ inside the outer horizon

Note that $P_0 = (\rho_0, \varphi_0, 0)$ and the null-geodesics γ_-, γ_+ starting at P_0 are lying in the equatorial plane $z = 0$, but U_0 is a neighborhood of P_0 in (ρ, φ, z) space. Therefore the null-geodesics starting on $\operatorname{supp}\chi_0$ are lying not only in the equatorial plane and the solutions $u^\pm$ constructed in §2 are the solutions of the equation (2.7) in the three-dimensional space.

Theorem 8.2. *Both u_0^+ and u_0^- cross the outer and the inner horizons and both end on the singularity ring $(\rho - a)^2 + z^2 = 0$.*

Proof: To prove this we shall show that all "plus" and "minus" null-geodesics starting on $\operatorname{supp}\chi_0$ will end on $(\rho - a)^2 + z^2 = 0$. Thus we need to estimate the behavior of null-geodesics in $\mathbb{R}^3$ starting on $\operatorname{supp}\chi_0$ and this is more difficult than the estimates for the null-geodesics lying in the equatorial plane.

We shall start with u_0^+ and will describe the behavior of plus null-geodesics $\gamma_+^{(1)}$ starting on $\operatorname{supp}\chi_0$ and close to γ^+, i.e. $\gamma_+^{(1)}$ is the projecion on (ρ, φ, z) of the null-bicharacteristic, with the initial data $(\rho', \varphi', z', \eta_\rho, \eta_\varphi, \eta_z)$ where (ρ', φ', z') are cloze to $(\rho_0, \varphi_0, 0)$ and $(\eta_\rho, \eta_\varphi, \eta_z)$ are the same as for γ^+. Note that z' is small but not equal to zero. Suppose, for the definiteness, that $z' > 0$.

We have the same formulas as in §7 and §8.1, with the variables z, ξ_z added:

$$\frac{d\rho^+}{dx_0} = -\frac{\sqrt{\Delta_2^+}}{\sqrt{\Delta_1^+}}, \quad \frac{dz}{dx_0} = -\frac{\sqrt{\Delta_3^+}}{\sqrt{\Delta_1^+}}, \tag{8.12}$$

where

$$\Delta_1^+ = (1 + K)\Big(\xi_\rho^2 + \xi_z^2 + \frac{\xi_\varphi^2}{\rho^2}\Big) - K\Big(b_\rho\xi_\rho + b_z\xi_z + b_\varphi\frac{\xi_\varphi}{\rho}\Big)^2. \tag{8.13}$$

Now $K = \frac{2mr^3}{r^4 + a^2z^2}$ where r is defined as in (1.5). Also

$$\Delta_2^+ = K(-\xi_0^+ + b_z\xi_z + b_\varphi\frac{\xi_\varphi}{\rho})^2 - (Kb_\rho^2 - 1)\Big((\xi_0^+)^2 - \xi_z^2 - \frac{\xi_\varphi^2}{\rho^2}\Big), \tag{8.14}$$

$$\Delta_3^+ = K(-\xi_0^+ + b_\rho\xi_\rho + b_\varphi\frac{\xi_\varphi}{\rho})^2 + (Kb_z^2 - 1)\Big((\xi_0^+)^2 - \xi_\rho^2 - \frac{\xi_\varphi^2}{\rho^2}\Big). \tag{8.15}$$

Near the singularity ring $(\rho - a)^2 + z^2 = 0$ we have from (1.5)
(8.16)

$$C_1'((|\rho{-}a|{+}|z|) \le C_1\sqrt{(\rho - a)^2 + z^2} \le r^2 \le C_2\sqrt{(\rho - a)^2 + z^2} \le C_2(|\rho{-}a|{+}|z|),$$

i.e. $r \sim (|\rho - a| + |z|)^{\frac{1}{2}}$ where $\alpha \sim \beta$ means $C_1\beta \le \alpha \le C_2\beta$.

Therefore from (1.4) and (8.16) we get

$$K \sim \frac{(|\rho - a| + |z|)^{\frac{3}{2}}}{(|\rho - a| + |z|)^2 + a^2 z^2} \sim \frac{1}{(|\rho - a| + |z|)^{\frac{1}{2}}}, \tag{8.17}$$

$$\begin{gathered} b_\rho = \frac{r\rho}{r^2 + a^2} \sim (|\rho - a| + |z|)^{\frac{1}{2}}, \quad b_z = \frac{z}{r} \sim \frac{z}{(|\rho - a| + |z|)^{\frac{1}{2}}}, \\ \frac{\xi_\varphi}{\rho} = \frac{a}{\rho} \sim 1, \quad b_\varphi = \frac{a\rho}{r^2 + a^2} \sim 1, \quad b_\varphi \frac{\xi_\varphi}{\rho} = \frac{a^2}{r^2 + a^2} \sim 1 - O(r^2). \end{gathered} \tag{8.18}$$

We shall estimate ξ_ρ^2 and ξ_z^2 from above. It follos from (1.2) with $H = 0$ that

$$\xi_\rho^2 + \xi_z^2 = K\Big(b_\rho\xi_\rho + b_z\xi_z - \xi_0^+ + b_\varphi\frac{\xi_\varphi}{\rho}\Big)^2 + (\xi_0^+)^2 - \frac{\xi_\varphi^2}{\rho^2}. \tag{8.19}$$

Therefore

$$\begin{aligned} \xi_\rho^2 + \xi_z^2 \le K\Big(- \xi_0^+ + b_\varphi\frac{a}{\rho} + b_z\xi_z\Big)^2 + 2K\Big| - \xi_0^+ + b_\varphi\frac{a}{\rho} + b_z\xi_z\Big| b_\rho|\xi_\rho| \\ + K b_\rho^2|\xi_\rho|^2 + \frac{\rho^2 - a^2}{\rho^2}. \end{aligned}$$

Using the identity $2cd = \varepsilon^2c^2 + \frac{d^2}{\varepsilon^2} - \big(\varepsilon c - \frac{d}{\varepsilon}\big)^2$ and choosing $\varepsilon^2 = \frac{Kb_\rho^2}{1 - Kb_\rho^2}$, $d = \sqrt{K}b_\rho|\xi_\rho|$, $c = \sqrt{K}\big| - \xi_0^+ + b_\varphi\frac{a}{\rho} + b_z\xi_z\big|$, we get

$$\begin{aligned} \xi_\rho^2 + \xi_z^2 \le \frac{1}{1 - Kb_\rho^2}K\Big(- \xi_0^+ + b_\varphi\frac{a}{\rho} + b_z\xi_z\Big)^2 + \xi_\rho^2 + \frac{\rho^2 - a^2}{\rho^2} \\ - \Big|\varepsilon\sqrt{K}\Big| - \xi_0^+ + b_\varphi\frac{a}{\rho} + b_z\xi_z\Big| - \frac{\sqrt{K}}{\varepsilon}b_\rho|\xi_\rho|\Big|^2. \end{aligned} \tag{8.20}$$

Canceling $|\xi_\rho^2|$ in (8.20) we get

$$|\xi_z|^2 \le \frac{1}{1 - Kb_\varphi^2}K\Big(- \xi_0^+ + b_\varphi\frac{a}{\rho} + b_z\xi_z\Big)^2 + \frac{\rho^2 - a^2}{\rho^2} - I_1^2, \tag{8.21}$$

where

$$(8.22)\qquad I_1 = |\varepsilon\sqrt{K}| - \xi_0^+ + b_\varphi \frac{a}{\rho} + b_z\xi_z| - \frac{\sqrt{K}}{\varepsilon} b_\rho|\xi_\rho|\,\big|,\quad \varepsilon^2 = \frac{Kb_\rho^2}{1-Kb_\rho^2}.$$

Using (8.21) we have

$$(8.23)$$

$$\begin{aligned}
&\Delta_2^+ = K\Big(-\xi_0^+ + b_\varphi\frac{a}{\rho} + b_z\xi_z\Big)^2 + (1-Kb_\rho^2)\Big(\frac{\rho^2-a^2}{\rho^2} - \xi_z^2\Big)\\
&\geq K\Big(-\xi_0^+ + b_\varphi\frac{a}{\rho} + b_z\xi_z\Big)^2\\
&+ (1-Kb_\rho^2)\Big(\frac{\rho^2-a^2}{\rho^2} - \frac{1}{1-Kb_\rho^2}K\Big(-\xi_0^+ + b_\varphi\frac{a}{\rho} + b_z\xi_z\Big)^2 - \frac{\rho^2-a^2}{\rho^2} + I_1^2\Big)\\
&= (1-Kb_\rho^2)I_1^2 \geq I_1^2.
\end{aligned}$$

Analogously we have, changing b_ρ to b_z and $|\xi_\rho|$ to $|\xi_z|$

$$(8.24)\qquad \Delta_3^+ \geq (1-Kb_z^2)I_2^2 \geq I_2^2,$$

where

$$(8.25)\qquad I_2 = \big|\varepsilon_1\sqrt{K}\big| - \xi_0^+ + b_\varphi\frac{a}{\rho} + b_\rho\xi_\rho| - \frac{\sqrt{K}}{\varepsilon_1}|b_z|\,|\xi_z|\big|,$$

where $\varepsilon_1 = \frac{Kb_z^2}{1-Kb_z^2}$.

Now estimate Δ_1^+. We have

$$\Delta_1^+ = (1+K)\Big(\xi_\rho^2 + \xi_z^2 + \frac{a^2}{\rho^2}\Big) - K\Big(b_\rho\xi_\rho + b_z\xi_z + b_\varphi\frac{a}{\rho}\Big)^2.$$

It follows from (8.19) that

$$\begin{aligned}
(8.26)\qquad \xi_\rho^2 + \xi_z^2 &\leq 2K\Big(-\xi_0^+ + b_\varphi\frac{a}{\rho}\Big)^2 + 2K(b_\rho\xi_\rho + b_z\xi_z)^2 + \frac{\rho^2-a^2}{\rho^2}\\
&\leq 2K\Big(-1 + \frac{a^2}{r^2+a^2}\Big)^2 + 2K(b_\rho^2 + b_z^2)(\xi_\rho^2 + \xi_z^2) + \frac{\rho^2-a^2}{\rho^2}.
\end{aligned}$$

Since $b_\rho^2 + b_z^2 = 1 - b_\varphi^2 = 1 - \frac{a^2\rho^2}{(r^2+a^2)^2} = O(r^2)$ we get from (8.17), (8.18)

$$(8.27)\ \ \xi_\rho^2 + \xi_z^2 \leq (1-2K(b_\rho^2+b_z^2))^{-1}\Big(\frac{\rho^2-a^2}{\rho^2} + C\delta^{\frac{3}{2}}\Big) \leq C_1\Big(\frac{\rho^2-a^2}{\rho^2} + C\delta^{\frac{3}{2}}\Big),$$

where

$$\delta = |\rho - a| + |z|. \tag{8.28}$$

Since $K\frac{a^2}{\rho^2} - Kb_\varphi^2\frac{a^2}{\rho^2} = O(\delta^{\frac{1}{2}})$, we get, using (8.27)

$$1 - C\delta^{\frac{1}{2}} < \Delta_1^+ < 1 + C\delta^{\frac{1}{2}}. \tag{8.29}$$

We will need also the estimate of $\xi_\rho^2 + \xi_z^2$ from below. It follows from (8.19), using the inequality of the form $(c+d)^2 \geq \frac{c^2}{2} - d^2$, that

$$\begin{aligned}\xi_\rho^2 + \xi_z^2 &\geq \frac{K}{2}\Big(- \xi_0^+ + b_\varphi\frac{a}{\rho}\Big)^2 - K(b_\rho\xi_\rho + b_z\xi_z)^2 + \frac{\rho^2 - a^2}{\rho^2} \\ &\geq \frac{K}{2}\Big(- \xi_0^+ + b_\varphi\frac{a}{\rho}\Big)^2 - K(b_\rho^2 + b_z^2)(\xi_\rho^2 + \xi_2^2) + \frac{\rho^2 - a^2}{\rho^2}.\end{aligned}$$

Therefore

$$\xi_\rho^2 + \xi_2^2 \geq \Big(1 + K(b_\rho^2 + b_z^2)\Big)^{-1}\Big(\frac{K}{2}\big(- \xi_0^+ + b_\varphi\frac{a}{\rho}\big)^2 + \frac{\rho^2 - a^2}{\rho^2}\Big). \tag{8.30}$$

Using (8.17), (8.18) we got

$$\xi_\rho^2 + \xi_z^2 \geq C\big(|\rho - a| + \delta^{\frac{3}{2}}\big) \geq C\delta^{\frac{3}{2}}. \tag{8.31}$$

Let $(\rho', \varphi', z', \eta_\rho, \eta_\varphi, \eta_z)$ be the initial data for the null-bicharacteristic system (2.1). Assuming that there are no focal points, there is a one-to-one correspondence between $(\rho(x_0), z(x_0))$ and (ρ', z'). Therefore $z(x_0) > 0$ if $z(0) = z' > 0$. Hence we shall write $\delta(x_0) = \rho(x_0) - a + z(x_0)$ instead of $\delta = |\rho(x_0) - \alpha| + |z|$ since $z(x_0) > 0$ and $\rho(x_0) - a > 0$.

We have from (8.12), (8.23), (8.24), (8.29) that

$$\frac{d\delta}{dx_0} = \frac{d\rho}{dx_0} + \frac{dz}{dx_0} = -\frac{\sqrt{\Delta_2^+}}{\sqrt{\Delta_1^+}} - \frac{\sqrt{\Delta_3^+}}{\sqrt{\Delta_1^+}} \leq -C(I_1 + I_2), \tag{8.32}$$

where $\delta = \rho - a + z$.

Note that (cf. (8.17), (8.18))

$$\begin{aligned}I_1 = &\Big|\frac{K^{\frac{1}{2}}b_\rho K^{\frac{1}{2}}}{(1 - Kb_\rho^2)^{\frac{1}{2}}}\big| - \xi_0^+ + b_\varphi\frac{a}{\rho} + b_z\xi_z\big| - \frac{\sqrt{K}(1 - Kb_\rho^2)^{\frac{1}{2}}b_\rho|\xi_\rho|}{\sqrt{K}|b_\rho|}\Big| \\ \geq &|1 - Kb_\rho^2|^{\frac{1}{2}}|\xi_\rho| - C_1\delta \geq C|\xi_\rho| - C_1\delta.\end{aligned} \tag{8.33}$$

Analogously

$$I_2 \geq C|\xi_z| - C_2\delta. \tag{8.34}$$

Using the estimate (8.31) we get

$$\frac{d\delta}{dx_0} \leq -C(I_1+I_2) \leq -C(|\xi_\rho|+|\xi_z|+C_3\delta) \leq -C\delta^{\frac{3}{4}}+C_1\delta \leq -C_2\delta^{\frac{3}{4}}. \tag{8.35}$$

As in (8.6) we conclude from (8.35) that $\delta(t_0)=0$ for some t_0. Therefore all null-geodesics $\gamma_+^{(1)}$ end on the singularity ring $(\rho-a)^2+z^2=0$.

Now we shall study the behavior of null-geodesics $\gamma_-^{(1)}$ close to γ^- inside the outer event horizon. Note that the case of γ^- null-geodesics differ from the case of γ^+ null-geodesics only by $\xi_0^- = \frac{K_0-1}{K_0+1}$ replacing $\xi_0^+=1$. Here K_0 is the $K=\frac{2mr^3}{r^4+a^2z^2}$ at the initial point $(\rho_0,\varphi_0,0)$. In particular, instead of (8.23), (8.24) we have

$$\Delta_2^- \geq I_3^2, \quad \Delta_3^- \geq I_4^2, \tag{8.36}$$

where

$$\begin{aligned} I_3 &= \Big|\varepsilon\sqrt{K}| - \xi_0^- + b_\varphi\frac{a}{\rho} + b_z\xi_z| - \frac{\sqrt{K}}{\varepsilon}|b_\rho|\,|\xi_\rho|\Big|, \quad \varepsilon^2 = \frac{Kb_\rho^2}{1-Kb_\rho^2}, \\ I_4 &= \Big|\varepsilon_1\sqrt{K}| - \xi_0^- + b_\varphi\frac{a}{\rho} + b_\rho\xi_\rho| - \frac{\sqrt{K}}{\varepsilon_1}|b_z|\,|\xi_z|\Big|, \quad \varepsilon_1^2 = \frac{Kb_z^2}{1-Kb_z^2}. \end{aligned} \tag{8.37}$$

Estimate (8.27) takes the form

$$\xi_\rho^2+\xi_z^2 \leq CK\Big(-\xi_0^- + \frac{a^2}{r^2+a^2}\Big)^2 + C\Big|-\xi_0^- + \frac{a^2}{\rho^2}\Big| \leq C\delta^{-\frac{1}{2}}.. \tag{8.38}$$

The estimate of $\xi_\rho^2+\xi_z^2$ from below have the form (cf. (8.30))

$$\xi_\rho^2+\xi_z^2 \geq \frac{C}{\delta^{\frac{1}{2}}}. \tag{8.39}$$

Combining (8.38) and (8.39) we get instead of (8.29)

$$C_1\delta^{-1} \leq \Delta_1^- \leq C_2\delta^{-1}. \tag{8.40}$$

Therefore it follows from (8.32), (8.40)

$$\frac{d\delta}{dx_0} = \frac{d\rho}{dx_0} + \frac{dz}{dx_0} \leq -\frac{C(I_3+I_4)}{\delta^{-\frac{1}{2}}}. \tag{8.41}$$

We have similarly to (8.33)

$$I_3 \geq C|\xi_\rho| - C_1, \quad I_4 \geq C|\xi_z| - C_1. \tag{8.42}$$

Therefore

$$\frac{d\delta}{dx_0} \leq -C\delta^{\frac{1}{2}}(|\xi_\rho| + |\xi_z| - C_1) \leq -C\delta^{\frac{1}{2}}\delta^{-\frac{1}{4}} = -C\delta^{\frac{1}{4}}. \tag{8.43}$$

From (8.43) analogously to (8.11) we get that $\delta = \rho^- - a + z^-$ vanish at the singularity ring. Therefore all null-geodesics $\gamma_-^{(1)}$ end on the singularity ring (see Fig. 1).

Hence both u_0^+ and u_0^- end on the singularity ring at a finite time.

9 The superradiance in the case of extremal ($\mathbf{a^2 = m^2}$) Kerr metric and naked singularity ($\mathbf{a^2 > m^2}$) Kerr matric

In both cases the construction of the solutions u^+ and u^- remains the same and we only indicate the difference in the behavior of the null-geodesics γ_+ and γ_-.

In the case of extremal metric $a^2 = m^2$ we have $r_+ = r_- = m$, i.e the outer and the inner horizons coincide. As in the case $a < m$ $\rho^+(x_0)$ and $\rho^-(x_0)$ cross the event horizon when x_0 increases.

Consider the behavior of $\rho^-(x_0)$ when x_0 decreases. Note that $Kb_\rho^2 - 1 = O((r-r_+)^2)$ and therefore (cf. (7.27)):

$$\xi_\rho^- \sim \frac{1}{(r-r_+)^2}. \tag{9.1}$$

Therefore $\sqrt{\Delta_1^-} \sim \frac{1}{(r-r_+)^2}$ and, similarly to (7.29),

$$\frac{d\rho^-}{dx_0} \leq C_0(\rho - \hat{\rho}_+)^2, \quad C_0 > 0, \tag{9.2}$$

where $\hat{\rho}_+ = \sqrt{a^2 + m^2}$.

Hence $-\frac{1}{\rho^- - \hat{\rho}_+} \leq C_0(x_0 + C_1)$, or $0 < \rho^-(x_0) - \hat{\rho}_+ \leq \frac{1}{-C_0(x_0+C_1)}$. Thus $\rho^-(x_0) \to \hat{\rho}_+$ when $x_0 \to -\infty$. Therefore the difference with nonextremal case $a < m$ is that $\rho^-(x_0) - \hat{\rho}_+$ decays as $O\big(\frac{1}{x_0}\big)$ in the extremal case and $\rho^-(x_0) - \hat{\rho}_+$ decays exponentially in the nonextremal case.

Consider the case when $a^2 > m^2$. Then the outer and the inner event horizons disappear and this case is called the case of naked singularity. Note that the ergoregion is $a < \rho < \sqrt{4m^2 + a^2}$.

Theorem 9.1. *In the case $a^2 > m^2$, i. e. in the case of the naked singularity, the null-geodesics γ^+ starts at a point P_0 inside ergosphere and reaches the singularity ring $\rho = a$ at some finite time $x_0^{(1)} > 0$. When $x_0 < 0$ γ^+ tends to the infinity when $x_0 \to -\infty$. The null-geodesics γ^- also starts at P_0 and ends at the ring singularity at a finite time $x_0^{(2)} > 0$. When $x_0 < 0$ γ^- first reaches a turning point ρ_* at $x^* < 0$ when x_0 decreases and then it turns and approaches the singularity at some finite time $x_0^{(3)} < x_0^*$ (see Fig. 2).*

Proof: Consider the behavior of $\rho^-(x_0)$ when $x_0 < 0$. As in (7.26) we can prove that there exists a turning point ρ_* where $a < \rho_0 < \rho_* < \sqrt{4m^2 + a^2}$ by showing that $\Delta_2^-(\rho_0) > 0$ and $\Delta_2^-(\sqrt{4m^2 + a^2} < 0$. Using that $\rho_0 > a$, we have $\xi_0^- < \frac{2m-a}{2m+a}$. Then $\frac{2m-a}{2m+a} < \frac{2a^2}{4m^2+a^2}$ when $a > m$ and this proves that $\Delta_2^-(\sqrt{4m^2 + a^2}) < 0$ (cf. (7.26)). Therefore there exists a turning point ρ_* where $\Delta_2^-(\rho_*) = 0$. Let $x_0^{(3)} < 0$ be such that $\rho(x_0^{(3)} = \rho_*$. When $x_0 < x_0^{(3)}$ the equation for $\rho^-(x_0)$ has the form $\frac{d\rho^-(x_0)}{dx_0} = \frac{\sqrt{\Delta_2^-}}{\sqrt{\Delta_1^-}}$ (cf. (7.27)). Note that $1 - Kb_\rho = \frac{(r-m)^2+a^2-m^2}{r^2+a^2} > \frac{a^2-m^2}{r^2+m^2}$. Therefore when $a^2 - m^2$ is small and r is close to m we have

$$\xi_\rho^- = O\Big(\frac{1}{a^2 - m^2}\Big). \tag{9.3}$$

Hence

$$\frac{d\varphi}{d\rho} = O\Big(\frac{1}{a^2 - m^2}\Big) \tag{9.4}$$

is large when $a^2 - m^2$ is small and r is close to m (cf. (7.29)). Therefore γ^- makes several rounds near $r = m$ and the number of these rounds becomes larger when $a^2 - m^2$ becomes smaller. When x_0 decreases further γ^- approaches the singularity $\rho = a$. Near $\rho = a$ we have, using (8.8) and (8.10), that (cf. (8.11))

$$\frac{d\rho^-}{dx_0} = \frac{\sqrt{\Delta_2^-}}{\sqrt{\Delta_1^-}} \leq \frac{C_0|\rho - a|^{-\frac{1}{4}}}{|\rho - a|^{-\frac{1}{2}}} = C_0|\rho - a|^{\frac{1}{4}}. \tag{9.5}$$

Similarly to (8.11) we have $\frac{4}{3}(\rho - a)^{\frac{3}{4}} \leq C_0 x_0 + C$. Therefore

$$\rho^-(x_0) - a \leq \frac{3}{4}C(x_0 - x_0^{(4)})^{\frac{4}{3}}, \quad x_0 > x_0^{(4)}, \tag{9.6}$$

i.e. $\rho(x_0^{(4)}) = a$. Also $\left|\frac{d\varphi}{d\rho}\right| \leq C(\rho - a)^{-\frac{1}{4}}$. Hence $|\varphi(x_0) - \varphi_0| \leq C_0|\rho - \alpha|^{\frac{3}{4}}$. Therefore the null-geodesics γ^- ends at the point (a, φ_0) when $x_0 \to x_0^{(2)}$ (see Fig. 2).

The behavior of γ^+ and γ^- for $x_0 > 0$ and γ^+ for $x_0 < 0$ is the same as in the case $a^2 < m^2$, and the proof is also the same.



Figure 2: The case $a > m$.
Null-geodesics γ_+ starts at P_0 and reaches $\rho = a$ at a finite time $x_0^{(1)} > 0$. When $x_0 \to -\infty$ γ_+ tends to infinity. Null-geodesics γ_- starts at P_0 and reaches $\rho = a$ at finite time $x_0^{(2)} > 0$. When $x_0 < 0$ γ_- passes a turning point, makes several rounds and reaches $\rho = a$ at some time $x_0^{(3)} < 0$.

References

[1] Brito K., Cardoso V., Pani P., Superradiance, Lecture notes in Physics, vol. 906, Springer-Verlag (2015)

[2] Dafermos, M., Rodnianski, I., Shlapentoch-Rothman, Y., A scattering theory for the wave equation on Kerr black hole exteriors, arXiv 1412.8379

[3] Eskin, G., Lectures on linear partial differential equations, GSM 123, AMS, Providence, RI (2011)

[4] Eskin, G., Inverse hyperbolic problems and optical black holes, Commun. Math. Phys. 297, 817-839 (2010)

[5] Eskin, G., Hall, M., Stationary black holes metrics in two space dimensions, Inverse Problems 32 (2016), 095006

[6] Frolov, V., Novikov, I, Black holes physics: basic concepts and new developments, Kluger Dordrecht (1998)

[7] Finster, F., Kamran, N., Smoller, J., Yau, S.-T., A rigorous treatment of energy extraction from a rotating black hole, Commun. Math. Phys. 287, 829-847 (2009)

[8] Gordon, W., Zur lightfortpflanzung nach der relativittstheorie (1923), Ann. Phys. (Leipzig), 12,421

[9] Leonhardt, U., Piwnicki, P., Relativistic effects of light in moving media with extremely low group velocity, Phys. Rev. Lett. (2000), 84, 822

[10] Maslov, V., Fedoryuk, M., Semi-classical approximation in quantum mechanics, Reidel, Dordrecht (1981)

[11] Penrose, R., Gravitational collapse: The role of general relativity, Rev. del Nuovo Cimento 1(1969) 252-276

[12] Starobinsky, A., Amplification of waves during reflection from a black hole, Soviet Physics JETP, 37 (1973), 28-32

[13] Visser, M., Acoustic black holes horizons, ergospheres and Hawking radiation, Class. Quant. Grav., 15 (6), 1767-1791 (1998)

[14] Visser, M., The Kerr space-time: A brief introduction, The Kerr space-time, Cambridge, University Press (2009)

[15] Wald, R., General Relativity, University of Chicago Press (1984)

[16] Zeldovich, Ya. Soviet Physics JETP 35 (1972), 1085-1087